Gastroenterology Referral Form





Fax completed form to: _____

| PATIENT INFORMATION | | | | | | | |
|--|--|---------------------------|-------------------------------|-----------------|------------------|------------------------|----------------------|
| Patient Name: | D | Date of Birth: | | | Referral Date: | | |
| Address: | City/State/Zip: | | | | | | |
| Home Phone: | Ce | ell Phone: | | | Work Phone: | | |
| Secondary Contact: | H | eight: | Weight: | | Male Fer | nale | |
| Patient Diagnosis & ICD-10: | | | | | | | |
| Allergies: | | | | | | | |
| PROVIDER INFORMATION | | | | | | | |
| Physician Name: | Li | ic.#: | | DEA#: | | | |
| Practice Name: | | | NPI#: | | | | |
| Address: | | | | City/State/Zip: | | | |
| Office Contact: | P | hone: | | | Fax: | | |
| Supervisory Physician (if applicable): | | | | | | | |
| PLEASE ATTACH | | | | | | | |
| Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable Vaccine status (any vaccination) and documentation of any recent vaccinations TB lab results within last 12 months HBV lab results within last 12 months (Infliximabs only) Liver enzymes lab results (Skyrizi only) Bilirubin levels (Skyrizi only) Letter of medical necessity if drug dosing or indication is outside of FDA guidelines | | | | | | | |
| NURSING & LAB ORDERS | | | | | | | |
| Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 10units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line | | | | | | | |
| Lab Orders: Lab Date & Frequency: | | | | | | | |
| PRESCRIPTION ORDERS | | | | | | | |
| Anaphylaxis Kit: | Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125 | | | | | | 5mg IV as needed |
| (Check all that apply) Pre-Medications: | Diphenhydraminemg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IVminutes prior to infusion | | | | | | |
| | | | | | | | |
| | | | | | | | |
| Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary | | | | | | | |
| PRODUCT | PRESCRIPTION INFORMATION | | | | | | REFILLS |
| Is this a first dose? Y | es No If No, when was last dose given? | | nen is patient due for next d | | | T I | |
| CIMZIA® | 200x2 Prefilled Syringe | | subcutaneously at weeks 0, | , 2 and 4 | | 1 Kit | NONE |
| | 200x2 LYO Powder Inject 400mg subcutaneously once every 4 weeks 4 week supply | | | | | NONE | |
| ENTYVIO | Induction: 300mg IV infusion over 30 minutes at week 0, 2, and 6 Maintenance: 300mg IV infusion over 30 minutes every weeks | | | | | | NONE |
| | Or Prefilled Pen 108 mg SC every 2 weeks | | | | | 2 pens, 13 refills | |
| HUMIRA® Citrate Free | Crohn's/UC Starter Package (8mg - 80mg Pens) Inject 160mg given as Two 80mg SubQ Day 1 OR One 80mg SubQ Days 1 & 2, then Week 2 inject 80 mg subcutaneously on Day 15 Work 4 - Livingt 40mg gubgutaneously on growther week Week supply | | | | NONE | | |
| INFLIXIMAB | 40Hig Feir 40Hig FF3 Week 4+: Hiject 40Hig subcutatiously every other week | | | | | NONE | |
| Avsola | | | | | | | NONL |
| Inflectra | Inflectra (Note: Round to nearest 100mg for Medicaid patients) | | | | | | |
| Remicade | | | | | | | |
| Renflexis | If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert. | | | | | | |
| SKYRIZI | Induction: 600mg IV infusion over one hour at week 0, 4, and 8 | | | | | | NONE |
| | Maintenance: 360mg SC injection at Week 12, and every 8 weeks thereafter | | | | | | |
| | Induction (Adult Dosing–Based on body weight of patient at time of dosing): For patients 55kg or less administer 260mg IV infusion over at least 1 hour x 1 dose | | | | | | NONE |
| STELARA | For patients more than 55kg to 85kg administer 390mg IV infusion over at least 1 hour x 1 dose | | | | | | |
| 3122.001 | For patients more than 85kg administer 520mg IV infusion over at least 1 hour x 1 dose | | | | | | |
| | Maintenance: 90 mg SubQ injection weeks after induction and every weeks thereafter | | | | | | |
| OTHER | | | | | | | NONE |
| By signing this form a | ınd utilizing our services, you are authorizing E | ventus Rx to serve as you | r prior authorization desig | nated agent in | dealing with med | dical and prescription | insurance companies. |

Prescriber's Signature
<u>Dispense as Written</u>

Print Name

Date

Prescriber's Signature Substitution Permitted **Print Name**

Date

