Gastroenterology Referral Form an

Fax completed form to: 833-908-1122







PATIENT INFORMATION					
Patient Name:	Date	e of Birth:	Referral Date:		
Address:			City/State/Zip:		
Home Phone: Cell Phone:		Phone:	Work Phone:		
Secondary Contact:	Heig	ıht: Weight:		Male Female	
Patient Diagnosis & ICD-10:					
Allergies:					
PROVIDER INFORMATION					
Physician Name:	Lic.#	:	DEA #:		
Practice Name:			NPI#:		
Address:		City/State/Z	City/State/Zip:		
Office Contact:			Fax:		
Supervisory Physician (if applicable):					
PLEASE ATTACH					
Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable Vaccine status (any vaccination) and documentation of any recent vaccinations TB lab results within last 12 months HBV lab results (Skyrizi only) Liver enzymes lab results (Skyrizi only) Bilirubin levels (Skyrizi only) Letter of medical necessity if drug dosing or indication is outside of FDA quidelines					
NURSING & LAB ORDERS					
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: Nacl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL0R 10units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line					
Lab Orders: Lab Date & Frequency:					
PRESCRIPTION ORDERS					
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed					
Check all that apply) Diphenhydramine mg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other					
Pre-Medications:					
(Check all that apply)					
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary					
PRODUCT	-	RESCRIPTION INFORM	-		REFILLS
	es No If No, when was last dose given?	When is patient due			
	Induction: 300mg IV infusion via gravityOR pump over 30 minutes at week 0, 2, and 6				NONE
ENTYVIO	Maintenance: 300mg IV infusion via gravityOR pump over 30 minutes every weeks				HONE
INFLIXIMAB		· · · · · · · · · · · · · · · · · · ·	•	h	NONE
Avsola					
Inflectra	Maintenance:mg/kgmglV infusion via gravityOR pump over at least 2 hours everyweeks(Note: Round to nearest 100mg for Medicaid patients)				
Remicade					
Renflexis	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.				
SKYRIZI	Induction: 600mg IV infusion via gravity	OR pump over one hour at we	ek 0, 4, and 8		NONE
	Maintenance: 360mg SC injection at Week 12, and every 8 weeks thereafter				
	Induction (Adult Dosing -Based on body weigh				
STELARA	For patients 55kg or less administer 260mg IV infusion via gravity OR pump over at least 1 hour x 1 dose				
	For patients more than 55kg to 85kg administer 390mg IV infusion via gravityOR pump over at least 1 hour x 1 dose For patients more than 85kg administer 520mg IV infusion via gravityOR pump over at least 1 hour x 1 dose				
					NONE
	Maintenance: 90mg SubQ injection weeks after induction and every weeks thereafter				
OTHER					NONE
OTHER					
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.					

Prescriber's Signature Print Name Date Prescriber's Signature Print Name Date

Dispense as Written Substitution Permitted





