## LEMTRADA® Referral Form





## Fax completed form to:

|   | PATIENT I   | NFORMATION                                    | I  |  |                 |  |
|---|---|---|--|--|-----------------|--|
| Patient Name:   | Date of Birth:  |   |  | Referral Date:                                       |                 |  |
| Address:  |   |   | City/State/Zi  | ):   |                 |  |
| Home Phone:   | Cell Phone:   |   |  | Work Phone:  |                 |  |
| Secondary Contact:  | Height: \   | Weight:                                       |  | Male Female  |                 |  |
| Patient Diagnosis & ICD-10:   |   |   |  |  |                 |  |
| Allergies:  |   |   |  |  |                 |  |
| Dhuri dan Nama  |   | INFORMATIO                                    | [  |  |                 |  |
| Physician Name:   | Lic.#:  |   | DEA #:<br>NPI#:  |  |                 |  |
| Practice Name:<br>Address:  |   |   |  |  |                 |  |
| Office Contact:   |   |   |  | City/State/Zip:<br>Fax:                              |                 |  |
| Supervisory Physician (if applicable):  |   |   |  |  |                 |  |
|   | MS CLINI  | CAL DETAILS                                   |  |  | -               |  |
| Type of MS: Primary progressive multiple sclere   |   |   |  |  |                 |  |
| Ambulation status: Able to ambulate more th   |   |   | erc  |  |                 |  |
| <b>Relapse details:</b> Two or more relapses within t   |   |   |  |  |                 |  |
| neupse details. Two of more relapses within t   |   | E ATTACH                                      |  |  | -               |  |
| Patient demographics & front/back copy of all in  |   |   | um creatinine  | evels urinalysis with cell counts urine protein to c | reatinine ratio |  |
| Abumatid from Abuma   |   |   |  |  |                 |  |
| Necent once visit notes, nistory & physical, iab & pertinent procedure results  |   |   |  |  |                 |  |
|   |   |   | accination) and documentation of any recent vaccinations |  |                 |  |
| Line access documentation/verification if applicable Letter of medical necessity if drug dosing or indication is outside of FDA guidelines  |   |   |  |  |                 |  |
|   | NURSING 8   | & LAB ORDERS                                  | S  |  | -               |  |
| Nurse Orders: Nurse to provide assessment, teaching   | ng, lab draws, medication administration and va   | scular access device inser                    | tion.  |  |                 |  |
| Flush Orders: Na/Cl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line |   |   |  |  |                 |  |
|   |   | 5/IIIL <b>UI</b> 100011                       | IC3/IIIL - J-JIIIL                                       |  |                 |  |
| <b>Oxygen:</b> Give 0 <sub>2</sub> at 2L/M per nasal cannula as need  |   |   |  |  |                 |  |
| Lab Orders: Lab Date & Frequency:   |   |   |  |  |                 |  |
|   | SUPPL   | Y ORDERS                                      |  |  |                 |  |
| Supply Orders: All supplies for vascular access line  | care, drug administration kit(s), pump, and IV pc   | ole will be provided as nec                   | essary   |  |                 |  |
| PRODUCT PRESCRIPTION INFORMATION  |   |   |  |  | REFILLS         |  |
|   |   |   |  |  |                 |  |
| Is this a first dose? Yes No If No, when w  | as last dose given?Whe  | en is patient due for next                    | dose?  |  |                 |  |
| Pre Meds: Hydroxyzine HCl 50mg po prior to start of infusion and every 6 hours prn #25  |   |   |  |  |                 |  |
|   | Acyclovir 200mg po BID for a minimum of 2 months or until CD4+ count is > or = to 200 cells per microliter, whichever occurs later #60 Refill: #1 |   |  |  |                 |  |
| Cetirizine 10mg po pric   |   |   | n 4mg po prn #   |  |                 |  |
| Promethazine 25mg p   | -   |   | 20mg prior to s  | tart of alemtuzumab infusion                         |                 |  |
|   | Acetaminophen 1000mg po prior to start of Lemtrada infusion and q6h prm Other:  |   |  |  |                 |  |
| LEMIKADA  | Note – If needed, please send pain prescription to retail pharmacy  |   |  |  |                 |  |
|   | Pre Infusion: Solu-Medrol 1000mg IV infusion in 500mL of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only                 |   |  |  |                 |  |
| Normal saline 0.9% 50   | Normal saline 0.9% 500ml IV prior to Lemtrada infusion on days 4 and 5  |   |  |  |                 |  |
| Initial Course: 12mg/   | Initial Course: 12mg/day IV infusion over 4 hours for 5 consecutive days  |   |  |  |                 |  |
| Subsequent Course:  | Subsequent Course: 12mg/day IV infusion over 4 hours for 3 consecutive days *To start at least 12 months after previous dose*                     |   |  |  |                 |  |
| Post Meds: Normal sa  | Post Meds: Normal saline 0.9% 500mL IV infusion over 1 hour post Lemtrada infusion  |   |  |  |                 |  |
| Ondansetron 4-8mg in 50-100mL 0.9% NaCl IV infusion over 15 minutes prn nausea  |   |   |  |  |                 |  |
|   | Epinephrine 1:1000 0.5-1mL IVP prn angioedema/hypotension/branchospasm/generalized urticaria  |   |  |  |                 |  |
| / SIDE EFFECT Kotorolac: 30mg IVP or  | Ketorolac: 30mg IVP over 3-5 minute   |   |  |  |                 |  |
|   | Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx 15 mins prn pruitis/rash  |   |  |  |                 |  |
| - Dipricingutatilite 301  |   |   |  |  |                 |  |
|   |   |   |  |  |                 |  |
| OTHER   |   | יישטע איז |  |  |                 |  |

Prescriber's Signature <u>Dispense as Written</u> Date

Prescriber's Signature Substitution Permitted

Print Name





Date