LEMTRADA® Referral Form

Fax completed form to: 833-908-1122





an amerita company

infusion solutions an amerita company

			company					
-	PATIENT INFORMATION							
Patient Name:	Date of Birth: Referral Date:							
Address:	City/State/Zip:							
Home Phone:	Cell Phone:	Work Phone:						
Secondary Contact:	Height: Weight:	Male Female						
Patient Diagnosis & ICD-10:								
Allergies:								
PROVIDER INFORMATION								
Physician Name:	Lic.#: DEA #:							
Practice Name:	NPI#:							
Address:	City/State/Zip:							
Office Contact:	Phone: Fax:							
Supervisory Physician (i								
	MS CLINICAL DETAILS							
Type of MS: Prima	nary progressive multiple sclerosis (PPMS) OR Relapsing multiple sclerosis (RMS)							
Ambulation status:	Able to ambulate more than 5 meters Able to ambulate without aid or rest for at least 100 meters							
Relapse details: T	Two or more relapses within the previous two years One relapse within the previous year							
•	PLEASE ATTACH							
Patient demographi		nine levels, urinalysis with cell counts, urine protein to ci	eatinine ratio					
5 1	notes, history & physical, lab & pertinent procedure results thyroid function tests							
	Pregnancy test results (if applicable	e)						
Current medication		d documentation of any recent vaccinations						
Line access docume	nentation/verification if applicable Letter of medical necessity if drug	dosing or indication is outside of FDA guidelines						
	NURSING & LAB ORDERS							
	to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/o							
Flush Orders: NaCl 0.99	1.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3	-5mL flush after post-infusion NS flush if indicated to m	aintain line					
Oxygen: Give 0 ₂ at 2L/I	L/M per nasal cannula as needed							
Lab Orders:	Lab Date & Frequency:							
SUPPLY ORDERS								
Supply Orders: All sup	upplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary							
PRODUCT	PRESCRIPTION INFORMATION	I	REFILLS					
Is this a first dose?								
IS UIIS & HISL UUSE!	Yes No If No, when was last dose given?When is patient due for next dose?							
	Pre Meds: Hydroxyzine HCI 50mg po prior to start of infusion and every 6 hours prn #25							
	Acyclovir 200mg po BID for a minimum of 2 months or until CD4+ count is > or = to 200 cells per microliter, whichever occurs later #60 Refill: #1							
	Cetirizine 10mg po prior to Lemtrada infusion Ondansetron 4mg po prn #25							
	Promethazine 25mg po prn #25 Famotidine 20mg prior to start of alemtuzumab infusion							
	Acetaminophen 1000mg po prior to start of Lemtrada infusion and q6h prn Other:							
	Note – If needed, please send pain prescription to retail pharmacy							
LEMTRADA	Pre Infusion: Solu-Medrol 1000mg IV infusion in 500mL of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only							
	Normal saline 0.9% 500ml IV prior to Lemtrada infusion on days 4 and 5							
	Initial Course: 12mg/day IV infusion via pumpOR gravity over 4 hours for 5 consecutive days							
	Subsequent Course: 12mg/day IV infusion via pumpOR gravity over 4 hours for 3 consecutive of	days *To start at least12 months after previous dose*						
	Post Meds: Normal saline 0.9% 500mL IV infusion over 1 hour post Lemtrada infusion							
ANAPHYLAXIS / Side effect	Ondansetron 4-8mg in 50-100mL 0.9% NaCl IV infusion over 15 minutes prn nausea							
	Epinephrine 1:1000 0.5-1mL IVP prn angioedema/hypotension/branchospasm/generalized urticaria							
	Ketorolac: 30mg IVP over 3-5 minute							
ORDERS								
	Ketorolac: 30mg IVP over 3-5 minute Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx 15 mins prn pruitis/rash							
ORDERS								

Prescriber's Signature	Print Name	Date	Prescriber's Signature	Print Name	Date
Dispense as Written			Substitution Permitted	\wedge	





