## **LEMTRADA®** Referral Form





## **Fax Completed Form To:**

**Phone:** 

|  | PATIEN  | NT INFORMATION                                |   |                  |  |
|--|---|---|---|------------------|--|
| Patient Name:  | Date of Birth:  | _   | Referral Date:  |                  |  |
| Address:   |   | City/State/                                   | Zip:  |                  |  |
| Home Phone:  | Cell Phone:   |   | Work Phone:   |                  |  |
| Secondary Contact:   | Height:   | Weight:                                       | Male Female   |                  |  |
| Patient Diagnosis & ICD  | -10:  |   |   |                  |  |
| Allergies:   |   |   |   |                  |  |
|  |   | DER INFORMATION                               |   |                  |  |
| Physician Name:  | Lic.#:  | DEA #:  |   |                  |  |
| Practice Name:   |   | NPI#:   |   |                  |  |
| Address:   | City/State/Zip:   |   |   |                  |  |
| Office Contact:  | Phone:  | Fax:  |   |                  |  |
| Supervisory Physician (i   |   | THICAL DETAILS                                |   |                  |  |
|  |   | LINICAL DETAILS                               |   |                  |  |
| Type of MS: Prima  | ry progressive multiple sclerosis (PPMS) <b>OR</b> Relapsing multipl  |   |   |                  |  |
| Ambulation status:   |   | ut aid or rest for at least 100 meters        |   |                  |  |
| Relapse details: T   | wo or more relapses within the previous two years One relapse w   | thin the previous year                        |   |                  |  |
|  | PL  | EASE ATTACH                                   |   |                  |  |
| Patient demographics & front/back copy of all insurance cards (prescription & medical)  CBC with differential, Serum creatinine levels, urinalysis with cell counts, urine protein to creatinine ratio |   |   |   |                  |  |
| Recent office visit no   | otes, history & physical, lab & pertinent procedure results   | thyroid function tests                        |   |                  |  |
|  | list & list of prior medications tried and failed (with dates)  |   | Pregnancy test results (if applicable)  |                  |  |
| ·  |   |   | Vaccine status (any vaccination) and documentation of any recent vaccinations |                  |  |
| Line access documentation/verification if applicable  Letter of medical necessity if drug dosing or indication is outside of FDA guidelines  |   |   |   |                  |  |
|  | NURSI   | NG & LAB ORDERS                               |   |                  |  |
| Nurse Orders: Nurse to   | provide assessment, teaching, lab draws, medication administration  | and vascular access device insertion and/or n | nanagement per physician orders.  |                  |  |
|  |   |   | nL flush after post-infusion NS flush if indicated to n                       | naintain line    |  |
|  |   | Touris, The Oil Touris, The 3 31              | ne nasiranci post iliusion ils nasirii iliatatea to il                        | idiritairi iiric |  |
| <del>-</del>   | M per nasal cannula as needed   |   |   |                  |  |
| Lab Orders:  |   | Lab Date & Frequency:                         |   |                  |  |
|  | SU  | PPLY ORDERS                                   |   |                  |  |
| Supply Orders: All sup   | plies for vascular access line care, drug administration kit(s), pump, an   | d IV pole will be provided as necessary       |   |                  |  |
| PRODUCT  |   | IPTION INFORMATION                            |   | REFILLS          |  |
|  | •   |   |   | REFILLS          |  |
| Is this a first dose?  | Yes No If No, when was last dose given?   | When is patient due for next dose?            |   |                  |  |
|  | Pre Meds: Hydroxyzine HCl 50mg po prior to start of infusion ar   | nd every 6 hours prn #25                      |   |                  |  |
| LEMTRADA  ANAPHYLAXIS / SIDE EFFECT  | Acyclovir 200mg po BID for a minimum of 2 months or until CD4+ count is > or = to 200 cells per microliter, whichever occurs later #60 Refill: #1 |   |   |                  |  |
|  | Cetirizine 10mg po prior to Lemtrada infusion   | Ondansetron 4mg po prn                        |   |                  |  |
|  | Promethazine 25mg po prn #25  | Famotidine 20mg prior to                      | start of alemtuzumab infusion   |                  |  |
|  | Acetaminophen 1000mg po prior to start of Lemtrada infusion a   |   |   |                  |  |
|  | Note — If needed, please send pain prescription to retail   | pharmacy                                      |   |                  |  |
|  | <b>Pre Infusion:</b> Solu-Medrol 1000mg IV infusion in 500mL of 0.9%<br>Normal saline 0.9% 500ml IV prior to Lemtrada infusion on days            |   | days 1, 2 and 3 only  |                  |  |
|  | Initial Course: 12mg/day IV infusion via pumpOR gravity over 4 hours for 5 consecutive days   |   |   |                  |  |
|  | Subsequent Course: 12mg/day IV infusion via pump OR gravity over 4 hours for 3 consecutive days *To start at least12 months after previous dose*  |   |   |                  |  |
|  | ,   |   | io start at reast 12 months arter previous dose                               |                  |  |
|  | Post Meds: Normal saline 0.9% 500mL IV infusion over 1 hour post Lemtrada infusion  |   |   |                  |  |
|  | Ondansetron 4-8mg in 50-100mL 0.9% NaCl IV infusion over 15 minutes prn nausea  |   |   |                  |  |
|  | Epinephrine 1:1000 0.5-1mL IVP prn angioedema/hypotension/branchospasm/generalized urticaria  |   |   |                  |  |
| / NIDE FEFF( I   |   | Ketorolac: 30mg IVP over 3-5 minute           |   |                  |  |
| ORDERS   | Ketorolac: 30mg IVP over 3-5 minute   |   |   | 1                |  |
|  | Ketorolac: 30mg IVP over 3-5 minute Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx   | 15 mins prn pruitis/rash                      |   |                  |  |
| ORDERS   |   | 15 mins prn pruitis/rash                      |   |                  |  |
| ORDERS<br>OTHER  | Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx   |   |   |                  |  |
| ORDERS<br>OTHER  |   |   | nt in dealing with medical and prescription insur                             | ance companies.  |  |

Prescriber's Signature Dispense as Written

**Print Name** 

Date

Prescriber's Signature **Substitution Permitted**  **Print Name** 

Date





