## Dermatology Referral Form

Phone:

Fax Completed Form To:



PATIENT INFORMATION						
Patient Name:		Date of Birth:		Referral Date:		
Address:		City/State/Zip:				
Home Phone:		Cell Phone:		Work Phone:		
Secondary Contact:		Height:	Weight:	Male Female		
Patient Diagnosis & ICD-	-10·	neight.	Treight.	marc remarc		
Allergies:						
PROVIDER INFORMATION						
Physician Name:		Lic.#:		DEA #:		
Practice Name:				NPI#:		
Address:				City/State/Zip:		
Office Contact:		Phone:		Fax:		
Supervisory Physician (if applicable):						
PLEASE ATTACH						
Patient demographics & front/back copy of all insurance cards (prescription & medical) TB lab results within last 12 months (Stelara, Simponi Aria, Ilumya & Infliximabs only)						
Recent office visit notes, history & physical, lab & pertinent procedure results HBV lab results within last 12 months (Infliximabs & Simponi Aria only)						
Current medication list & list of prior medications tried and failed (with dates) Letter of medical necessity if drug dosing or indication is outside of FDA guidelines						
NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.						
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line						
Lab Orders: Lab Date & Frequency:						
PRESCRIPTION ORDERS						
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed						
(Check all that apply) Diphenhydramine mg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other						
Pre-Medications: Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IVminutes prior to infusion						
(Check all that apply) Diphenhydramine mg PO OR IV infusion minutes prior to infusion Other						
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary						
PRODUCT   PRESCRIPTION INFORMATION   REFILI						
Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose?						
ILUMYA	100mg SC injection at 0 and 4 weeks then every 12 weeks					
INFLIXIMAB	Induction:mg/kg or	mg IV infusion via	gravity <b>OR</b> pump o	ver at least 2 hours at weeks 0, 2, and 6	NONE	
Avsola	Maintenance:mg/kg	mg IV infusion via	gravity <b>OR</b> pump of	over at least 2 hours every weeks		
Inflectra	(Note: Round to nearest 100mg for Medicaid patients)					
Remicade	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.					
Renflexis						
SIMPONI ARIA	2 mg/kg IV infusion via gravityOR pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter   000 mg IV infusion use 00 minutes Additional 000 mg IV infusion use 00 minutes are used a factorization of the second o					
SPEVIGO	900 mg IV infusion over 90 minutes Additional 900 mg IV infusion over 90 minutes one week after initial dose if flare symptoms persist					
	Psoriasis Adult Subcutaneous		۰ ۲. السبب ما الب ۲. من م	Durala		
STELARA	For patients <= 100 kg, 45 mg SC injection initially and 4 weeks later, followed by 45 mg every 12 weeks For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks					
	Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose)					
	For patients $<= 60 \text{ kg}$ , 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks					
	For patients 60 kg – 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks					
	For patients > 100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks					
	Psoriatic Arthritis Adult					
	45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks For patients with co-existent moderate-to-severe plaque psoriasis weighing >100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks					
XOLAIR	150 or 300 mg SC injection once every 4 weeks					
IG	For Immunoglobulin therapy please refer to Immunoglobulin Form					
OTHER		<u> </u>				
I	By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.					

Prescriber's Signature **Dispense as Written** 

**Print Name** 

Date

Prescriber's Signature **Substitution Permitted**  **Print Name** 

Date

