LEMTRADA® Referral Form



Fax Completed Form To:

Phone:

		DATIEN	T INFORMATION	т			
Patient Name:		Date of Birth:	I INFORMATION		Referral Date:		
Address:		City/State/Zip:					
Home Phone:		Cell Phone:		City/State/Li	Work Phone:		
Secondary Contact:		Height:	Weight:		Male Female		
Patient Diagnosis & ICI)-10·	Theight	reight		male remale		
Allergies:							
PROVIDER INFORMATION							
Physician Name:		Lic.#:		DEA #:			
Practice Name:		1		NPI#:			
Address:					n:		
Office Contact:	ce Contact: Phone:			City/State/Zip: Fax:			
Supervisory Physician (if applicable):							
MS CLINICAL DETAILS							
Type of MS: Primary progressive multiple sclerosis (PPMS) OR Relapsing multiple sclerosis (RMS)							
Ambulation status: Able to ambulate more than 5 meters Able to ambulate without aid or rest for at least 100 meters							
Relapse details: Two or more relapses within the previous two years One relapse within the previous year							
PLEASE ATTACH							
PLEASE AT FACH Patient demographics & front/back copy of all insurance cards (prescription & medical) CBC with differential, Serum creatinine levels, urinalysis with cell counts, urine protein to creatinine ratio							
thursd function tasts						reaurimerauo	
Recent office visit notes, history & physical, lab & pertinent procedure results Pregnancy test results (if applicable)							
Current medication	list & list of prior medications tried and failed			cumentation of any recent vac	cinations		
Line access documentation/verification if applicable Letter of medical necessity if drug dosing or indication is outside of FDA guidelines NURSING & LAB ORDERS							
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.							
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line							
Oxygen: Give O ₂ at 2L/M per nasal cannula as needed							
Lab Orders: Lab Date & Frequency:							
SUPPLY ORDERS							
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary							
PRODUCT	PRESCRIPTION INFORMATION I						REFILLS
Is this a first dose? Yes No If No, when was last dose given?When is patient due for next dose?							
LEMTRADA	Pre Meds: Hydroxyzine HCl 50mg po pr	ior to start of infusion and	every 6 hours prn #25				
	Acyclovir 200mg po BID for a minimum of 2 months or until CD4+ count is > or = to 200 cells per microliter, whichever occurs later #60 Refill: #1						
	Cetirizine 10mg po prior to Lemtrada infusion Ondansetron 4mg po prn #25						
	Promethazine 25mg po prn #25 Famotidine 20mg prior to start of alemtuzumab infusion						
	Acetaminophen 1000mg po prior to start of Lemtrada infusion and q6h prn Other:						
	Note – If needed, please send pain prescription to retail pharmacy						
	Pre Infusion: Solu-Medrol 1000mg IV infusion in 500mL of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution on the Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution on the Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution on the Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Named Solution on the Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on the Named Solution on the Named Solution of 0.9% NaCl over 1 hour prior to Lemtrada infusion on the Named Solution on the Named						
	Normal saline 0.9% 500ml IV prior to Lemtrada infusion on days 4 and 5						
	Initial Course: 12mg/day IV infusion via pump OR gravity over 4 hours for 5 consecutive days						
	Subsequent Course: 12mg/day IV infusion via pump OR gravity over 4 hours for 3 consecutive days *To start at least 12 months after previous dose*						
	Post Meds: Normal saline 0.9% 500mL IV infusion over 1 hour post Lemtrada infusion						
ANAPHYLAXIS / SIDE EFFECT ORDERS	Ondansetron 4-8mg in 50-100mL 0.9% NaCl IV infusion over 15 minutes prn nausea						
	Epinephrine 1:1000 0.5-1mL IVP prn angioedema/hypotension/branchospasm/generalized urticaria						
	Ketorolac: 30mg IVP over 3-5 minute						
	Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx 15 mins prn pruitis/rash						
OTHER							
By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.							

Prescriber's Signature Substitution Permitted

ACHC

Print Name

Date

