

# Gastroenterology Referral Form



Fax Completed Form To:

Phone:

| PATIENT INFORMATION         |                |                 |             |
|-----------------------------|----------------|-----------------|-------------|
| Patient Name:               | Date of Birth: | Referral Date:  |             |
| Address:                    |                | City/State/Zip: |             |
| Home Phone:                 | Cell Phone:    | Work Phone:     |             |
| Secondary Contact:          | Height:        | Weight:         | Male Female |
| Patient Diagnosis & ICD-10: |                |                 |             |
| Allergies:                  |                |                 |             |

| PROVIDER INFORMATION                   |        |                 |
|--|--------|-----------------|
| Physician Name:                        | Lic.#: | DEA #:          |
| Practice Name:                         |        | NPI#:           |
| Address:                               |        | City/State/Zip: |
| Office Contact:                        | Phone: | Fax:            |
| Supervisory Physician (if applicable): |        |                 |

| PLEASE ATTACH  |   |
|--|---|
| Patient demographics & front/back copy of all insurance cards (prescription & medical)<br>Recent office visit notes, history & physical, lab & pertinent procedure results<br>Current medication list & list of prior medications tried and failed (with dates)<br>Line access documentation/verification if applicable<br>Vaccine status (any vaccination) and documentation of any recent vaccinations | TB lab results within last 12 months<br>HBV lab results within last 12 months ( <i>Infliximabs only</i> )<br>Liver enzymes lab results ( <i>Skyrizi only</i> )<br>Bilirubin levels ( <i>Skyrizi only</i> )<br>Letter of medical necessity if drug dosing or indication is outside of FDA guidelines |

| NURSING & LAB ORDERS   |  |
|--|--|
| <b>Nurse Orders:</b> Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.                        |  |
| <b>Flush Orders:</b> NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line |  |
| <b>Lab Orders:</b> <span style="float: right;"><b>Lab Date &amp; Frequency:</b></span>   |  |

| PRESCRIPTION ORDERS   |   |   |                                       |
|---|---|---|---------------------------------------|
| <b>Anaphylaxis Kit:</b>   | Epinephrine 0.3mg IM as needed  | Solu-cortef 250mg-500mg IV as needed                    | Solu-Medrol 60mg - 125mg IV as needed |
| (Check all that apply)  | Diphenhydramine _____ mg IV as needed                                   | NS Hydration 500 ml IV over 30 minutes as needed        | Other _____                           |
| <b>Pre-Medications:</b>   | Acetaminophen _____ mg PO _____ minutes prior to infusion               | Solu-Medrol _____ mg IV _____ minutes prior to infusion |                                       |
| (Check all that apply)  | Diphenhydramine _____ mg PO ---OR--- IV _____ minutes prior to infusion | Other _____   |                                       |
| <b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary |   |   |                                       |

| PRODUCT  | PRESCRIPTION INFORMATION  | REFILLS                             |
|--|---|-------------------------------------|
| Is this a first dose? Yes No If No, when was last dose given? _____ When is patient due for next dose? _____ |   |                                     |
| ENTYVIO  | <b>Induction:</b> 300mg IV infusion over 30 minutes at week 0 and 2<br><b>Maintenance:</b> 300mg IV infusion over 30 minutes every _____ weeks<br>--OR-- Prefilled Pen 108mg SC every 2 weeks starting at week 6  | NONE<br>_____<br>2 pens, 13 refills |
| INFLIXIMAB<br>Avsola<br>Inflectra<br>Remicade<br>Renflexis   | <b>Induction:</b> _____ mg/kg or _____ mg IV infusion via gravity ---OR--- pump over at least 2 hours at weeks 0, 2, and 6<br><b>Maintenance:</b> _____ mg/kg _____ mg IV infusion via gravity ---OR--- pump over at least 2 hours every _____ weeks<br>(Note: Round to nearest 100mg for Medicaid patients)<br>If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.  | NONE<br>_____<br>_____              |
| OMVOH  | <b>Induction:</b> 300mg IV infusion via gravity ---OR--- pump over 30 minutes at week 0, 4, and 8<br><b>Maintenance:</b> 200mg SC injection (given as two consecutive injections of 100 mg each) at Week 12, and every 4 weeks thereafter   | NONE<br>_____<br>_____              |
| SKYRIZI  | <b>Induction:</b> 600mg IV infusion via gravity ---OR--- pump over one hour at week 0, 4, and 8<br><b>Maintenance:</b> 360mg SC injection at Week 12, and every 8 weeks thereafter  | NONE<br>_____<br>_____              |
| STELARA  | <b>Induction (Adult Dosing -Based on body weight of patient at time of dosing):</b><br>For patients 55kg or less administer 260mg IV infusion via gravity ---OR--- pump over at least 1 hour x 1 dose<br>For patients more than 55kg to 85kg administer 390mg IV infusion via gravity ---OR--- pump over at least 1 hour x 1 dose<br>For patients more than 85kg administer 520mg IV infusion via gravity ---OR--- pump over at least 1 hour x 1 dose<br><b>Maintenance:</b> 90mg SubQ injection _____ weeks after induction and every _____ weeks thereafter | NONE<br>_____<br>_____              |
| OTHER  |   | NONE<br>_____<br>_____              |

By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature \_\_\_\_\_  
 Dispense as Written \_\_\_\_\_  
 Print Name \_\_\_\_\_ Date \_\_\_\_\_

Prescriber's Signature \_\_\_\_\_  
 Substitution Permitted \_\_\_\_\_  
 Print Name \_\_\_\_\_ Date \_\_\_\_\_

