Krystexxa® Order Form



6

Fax Completed Form To:

Phone:

PATIENT INFORMATION					
Data and		r	FORMATION		
Patient Name:		Date of Birth:	<i>c</i> : <i>i</i> c: .	Referral Date:	
Address:		Call Disease	City/State		
Home Phone:		Cell Phone:	:	Work Phone:	
Secondary Contact:	10.	Height: We	ight:	Male Female	
Patient Diagnosis & ICD-10: Allergies:					
PROVIDER INFORMATION					
Physician Name: Lic.#: DEA #:					
Practice Name: NPI#:					
Address:			City/State/Zip:		
Office Contact:	Phone:			Fax:	
Supervisory Physician (if applicable):					
PLEASE ATTACH					
Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) G6PD deficiency results Verification that patient has discontinued or plans to discontinue oral urate lowering medications			Evidence of patient on concurrent immunomodulation therapy such as: methotrexate, mycophenolate, leflunomide, azathioprine, or cyclosporine (Evidence supports the combination of Krystexxa and an immunomodulator in improving the patient's response to therapy; consider adding an immunomodulator if clinically appropriate.) Baseline serum Uric Acid lab results Letter of medical necessity if drug dosing or indication is outside of FDA guidelines		
NURSING & LAB ORDERS					
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency:					
PRESCRIPTION ORDERS					
Anaphylaxis Kit:Epinephrine 0.3mg IM as neededSolu-cortef 250mg-500mg IV infusion as neededSolu-Medrol 60mg - 125mg IV infusion as needed(Check all that apply)Diphenhydramine mg IV infusion as neededNS Hydration 500 ml IV infusion over 30 minutes as neededOther					
Pre-Medications: Per prescribing information: Pre-medications of antihistamine, corticosteroid and analgesic is recommended (Check all that apply) Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IV infusionminutes prior to infusion Diphenhydraminemg POOR IV infusionminutes prior to infusion Other					
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary					
PRODUCT		PRESCRIPTION	INFORMATION		REFILLS
Is this a first dose?	Yes No If No, when was last dose given	?When	is patient due for next dose?		
8mg IV infusion via gravity or pump over at least 2 hours every 2 weeks Krystexxa Image: After first infusion, patient to have sUA level performed within 48 hours prior to each infusion. For KVO: NS 100mL via IV infusion over 1 hour. If sUA is < 6mg/dL, proceed.					
OTHER					
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.					

Prescriber's Signature **Dispense as Written**

Date

Prescriber's Signature **Substitution Permitted**

Print Name



Date