## REZZAYO® (rezafungin) Referral Form







Fax Completed Form To:

Phone:

PATIENT INFORMATION						
Patient Name:	Date of Birth:			Referral Date:		
Address:			City/State/Z		ip:	
Home Phone:		Cell Phone:			Work Phone:	
Secondary Contact:		Height: Weight:			Male Female	
Patient Diagnosis & ICD-10:						
Allergies:						
PROVIDER INFORMATION						
Physician Name:		Lic.#:		DEA#:		
Practice Name:				NPI#:		
Address:				City/State/Zip:		
Office Contact:		Phone:		Fax:		
Supervisory Physician (if applicable):						
PLEASE ATTACH						
Patient demographics & front/back copy of all insurance cards (prescription & medical)  Most recent liver function panel						
Recent office visit notes, history & physical, lab & pertinent procedure results  Culture & sensitivity results						
Current medication list & list of prior medications tried and failed (with dates)  Letter of medical necessity if drug dosing or indication is outside of FDA guidelines						
Line access documentation/verification if applicable  NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.  Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 10units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line						
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line  Lab Orders:						
Lab Date & Frequency:						
PRODUCT	PRESCRIPTION INFORMATION					REFILLS
Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose?						
Anaphylaxis Kit:	Epinephrine 0.3mg IM as needed Solu-Cortef 250mg-500mg IV infusion as needed Solu-Medrol 60mg - 125mg IV infusion as needed					
(Check all that apply)	Diphenhydramine mg IV infusi	<u> </u>	dration 500 ml IV infusion ove	r 30 minutes a	s needed Other	NONE
REZZAYO	Induction: 400mg IV in 250ml NS/D5W over 1 hour via gravity <b>OR</b> pump					NONE
	Maintenance: 200mg IV in 250ml NS/D5W over 1 hour via gravity OR pump once weekly beginning on day 8 for up to 4 doses					
OTHER DOSING REGIMEN						
By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.						
Prescriber's Signature	Print Name	Date	Prescriber's Signa	ture	Print Name D	ate





