

Krystexxa® Order Form

Fax Completed Form To:

Phone:



| PATIENT INFORMATION | | | |
|-----------------------------|----------------|-----------------|-------------|
| Patient Name: | Date of Birth: | Referral Date: | |
| Address: | | City/State/Zip: | |
| Home Phone: | Cell Phone: | Work Phone: | |
| Secondary Contact: | Height: | Weight: | Male Female |
| Patient Diagnosis & ICD-10: | | | |
| Allergies: | | | |

| PROVIDER INFORMATION | | |
|--|--------|-----------------|
| Physician Name: | Lic.#: | DEA #: |
| Practice Name: | | NPI#: |
| Address: | | City/State/Zip: |
| Office Contact: | Phone: | Fax: |
| Supervisory Physician (if applicable): | | |

| PLEASE ATTACH | |
|--|--|
| Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) G6PD deficiency results Verification that patient has discontinued or plans to discontinue oral urate lowering medications | Evidence of patient on concurrent immunomodulation therapy such as: methotrexate, mycophenolate, leflunomide, azathioprine, or cyclosporine (Evidence supports the combination of Krystexxa and an immunomodulator in improving the patient's response to therapy; consider adding an immunomodulator if clinically appropriate.) Baseline serum Uric Acid lab results Letter of medical necessity if drug dosing or indication is outside of FDA guidelines |

| NURSING & LAB ORDERS |
|---|
| Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency: |

| PRESCRIPTION ORDERS | | | |
|-------------------------|--|--|--|
| Anaphylaxis Kit: | Epinephrine 0.3mg IM as needed | Solu-cortef 250mg-500mg IV infusion as needed | Solu-Medrol 60mg - 125mg IV infusion as needed |
| (Check all that apply) | Diphenhydramine _____ mg IV infusion as needed | NS Hydration 500 ml IV infusion over 30 minutes as needed | Other _____ |
| Pre-Medications: | Per prescribing information: Pre-medications of antihistamine, corticosteroid and analgesic is recommended | | |
| (Check all that apply) | Acetaminophen _____ mg PO _____ minutes prior to infusion | Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion | |
| | Diphenhydramine _____ mg PO ---OR--- IV infusion _____ minutes prior to infusion | Other _____ | |

Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary

| PRODUCT | PRESCRIPTION INFORMATION | REFILLS |
|------------------------------|---|---------|
| Is this a first dose? Yes No | If No, when was last dose given? _____ When is patient due for next dose? _____ | |
| Krystexxa | 8mg IV infusion via gravity or pump over at least 2 hours every 2 weeks <input checked="" type="checkbox"/> After first infusion, patient to have sUA level performed within 48 hours prior to each infusion. For KVO: NS 100mL via IV infusion over 1 hour. If sUA is ≤ 6mg/dL, proceed. If sUA is > 6mg/dL, hold & contact provider. | _____ |
| OTHER | | _____ |

By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature _____
 Dispense as Written _____
 Print Name _____
 Date _____

Prescriber's Signature _____
 Substitution Permitted _____
 Print Name _____
 Date _____

