## Krystexxa® Order Form

Fax Completed Form To:

Phone:



PATIENT INFORMATION						
Patient Name:		Date of Birth:			Referral Date:	
Address:				City/State/Zip		
Home Phone:	Cell Phone:				Work Phone:	
Secondary Contact:		Height: Wei	aht:		Male Female	
Patient Diagnosis & ICD	-10:	<b>,</b>				
Allergies:						
PROVIDER INFORMATION						
Physician Name:	nysician Name: Lic.#:			DEA #:		
Practice Name:			NPI#:			
Address:				City/State/Zip:		
Office Contact:	Phone:			Fax:		
Supervisory Physician (if applicable):						
PLEASE ATTACH						
Patient demographics & front/back copy of all insurance cards (prescription & medical) Evidence of patient on concurrent immunomodulation therapy such as: methotrexate,						
musanhanalata laflunamida azathianzina az sudasnazina (Evidansa						
Recent office visit notes, history & physical, hab & per then procedure results				combination of Krystexxa and an immunomodulator in improving the patient's response to		
				therapy; consider adding an immunomodulator if clinically appropriate.)		
G6PD deficiency results Baseline s				Baseline serum Uric Acid lab results		
			Lattor of modical	Letter of medical necessity if drug dosing or indication is outside of FDA guidelines		
NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.						
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line						
Lab Orders:						
Lab Date & Frequency:						
PRESCRIPTION ORDERS						
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV infusion as needed Solu-Medrol 60mg - 125mg IV infusion as needed						
(Check all that apply) Diphenhydramine mg IV infusion as needed NS Hydration 500 ml IV infusion over 30 minutes as needed Other						
<b>Pre-Medications:</b> Per prescribing information: Pre-medications of antihistamine, corticosteroid and analgesic is recommended						
(Check all that apply) Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IV infusionminutes prior to infusion						
Diphenhydramine mg PO OR IV infusion minutes prior to infusion Other						
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary						
PRODUCT PRESCRIPTION INFORMATION REFILLS						
Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose?						
15 this a hist auge:	-					
8mg IV infusion via gravity or pump over at least 2 hours every 2 weeks						
	☑ After first infusion, patient to have sUA level performed within 48 hours prior to each infusion.					
Krystexxa	For KVO: NS 100mL via IV infusion over 1 hour.					
	If sUA is $\leq$ 6mg/dL, <b>proceed</b> .					
	If sUA is > 6mg/dL, <b>hold &amp; contact pro</b>	vider.				
	•					
OTHER						
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.						

Prescriber's Signature <u>Dispense as Written</u> Date

Prescriber's Signature Substitution Permitted Print Name

ACHC





