BLINCYTO® Order Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION					
Patient Name:		Date of Birth:			Referral Date:
Address:				City/State/Zip:	
Home Phone:		Cell Phone:			Work Phone:
Secondary Contact:		Height:	Weight:		🗆 Male 🛛 Female
Patient Diagnosis & ICD	-10:				
Allergies:					
PROVIDER INFORMATION					
Physician Name:		Lic.#:		DEA #:	
Practice Name:				NPI#:	
Address:				City/State/Zip:	
Office Contact:		Phone:		Fax:	
Supervisory Physician (if applicable):					
PLEASE ATTACH					
 Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable 			 Vaccine status (any vaccination) and documentation of any recent vaccinations TB lab results within last 12 months HBV lab results within last 12 months (Infliximabs only) Letter of medical necessity if drug dosing or indication is outside of FDA guidelines 		
NURSING & LAB ORDERS					
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.					
Lab Orders: Lab Date & Frequency:					
PRESCRIPTION ORDERS					
Anaphylaxis Kit: D Epinephrine 0.3mg IM as needed NS Hydration 500 ml IV infusion over 30 minutes as needed					
(Check all that apply) Diphenhydramine mg IV infusion as needed Other					
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary					
PRODUCT PRESCRIPTION INFORMATION					
□ Blinatumomab (BLINCYTO®)	Maintenance Orders (Consolidation cycles): Dispense up to 9 cycles as ordered. Current cycle number: Date current cycle initiated: Infuse 28 mcg/day IV infusion continuously via ambulatory pump (patient weight ≥ 45 kg) x 28 days, followed byday treatment-free interval. Infuse 15 mcg/m2/day IV infusion continuously via ambulatory pump. Medicare Orders: E0781 Ambulatory Infusion (1 per month), A4222 IV Admin Kit (1 per bag/cassette), A4221 IV supplies (1 per week) Ancillary Medication Orders: Patients Weighing ≥ 45 kg (Select one of the following): Dexamethasone 20 mg IV one hour before 1st dose of each new cycle (relapsed/refractory) or when restarting an infusion after an interruption of 4 or more hours in the first cycle. Prednisone 100mg IV infusion cont flux in between blinatumomab (Blincyto®) bag changes.] PliCC and Central Tunneled/Non-Tunneled: NS 5 mL pre-lab draw and 10 mL post-lab draw. For maintenance, heparin [(10 unit/mL) 5 mL or [(100 unit/mL) 3 mL every 24 hr to non-medication lumen. Implanted Port: When appropriate, NS 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance, heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance, heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance, heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance, heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance, heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance, heparin (100 unit/mL) 3 to 5 mL post-use at completion of cycle. For maintenance				
D OTHER					
By signing this form ar	nd utilizing our services, you are authorizing .	Amerita, Inc. to serve as	your prior authorization desi	ignated agent	t in dealing with medical and prescription insurance companies.

Prescriber's Signature Dispense as Written Date

Prescriber's Signature Substitution Permitted Print Name

Date





