

Dermatology Referral Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION			
Patient Name:	Date of Birth:	Referral Date:	
Address:	City/State/Zip:		
Home Phone:	Cell Phone:	Work Phone:	
Secondary Contact:	Height:	Weight:	<input type="checkbox"/> Male <input type="checkbox"/> Female
Patient Diagnosis & ICD-10:			
Allergies:			

PROVIDER INFORMATION			
Physician Name:	Lic.#:	DEA #:	
Practice Name:	NPI#:		
Address:	City/State/Zip:		
Office Contact:	Phone:	Fax:	
Supervisory Physician (if applicable):			

PLEASE ATTACH	
<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical)	<input type="checkbox"/> TB lab results within last 12 months (<i>Stelara, Simponi Aria, Ilumya & Infliximabs only</i>)
<input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results	<input type="checkbox"/> HBV lab results within last 12 months (<i>Infliximabs & Simponi Aria only</i>)
<input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates)	<input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guidelines

NURSING & LAB ORDERS	
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.	
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - <input type="checkbox"/> 10units/mL ---OR--- <input type="checkbox"/> 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line	
Lab Orders: Lab Date & Frequency:	

PRESCRIPTION ORDERS			
Anaphylaxis Kit:	<input type="checkbox"/> Epinephrine 0.3mg IM as needed	<input type="checkbox"/> Solu-cortef 250mg-500mg IV as needed	<input type="checkbox"/> Solu-Medrol 60mg - 125mg IV as needed
(Check all that apply)	<input type="checkbox"/> Diphenhydramine _____ mg IV as needed	<input type="checkbox"/> NS Hydration 500 ml IV over 30 minutes as needed	<input type="checkbox"/> Other _____
Pre-Medications:	<input type="checkbox"/> Acetaminophen _____ mg PO _____ minutes prior to infusion	<input type="checkbox"/> Solu-Medrol _____ mg IV _____ minutes prior to infusion	
(Check all that apply)	<input type="checkbox"/> Diphenhydramine _____ mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV infusion _____ minutes prior to infusion	<input type="checkbox"/> Other _____	
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No If No, when was last dose given? _____ When is patient due for next dose? _____		
<input type="checkbox"/> ILUMYA	100mg SC injection at 0 and 4 weeks then every 12 weeks	_____
<input type="checkbox"/> INFlixIMAB <input type="checkbox"/> Avsola <input type="checkbox"/> Inflectra <input type="checkbox"/> Remicade <input type="checkbox"/> Renflexis	<input type="checkbox"/> Induction: _____ mg/kg or _____ mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours at weeks 0, 2, and 6	NONE
	<input type="checkbox"/> Maintenance: _____ mg/kg _____ mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours every _____ weeks <i>(Note: Round to nearest 100mg for Medicaid patients)</i> If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.	_____
<input type="checkbox"/> SIMPONI ARIA	2 mg/kg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter	_____
<input type="checkbox"/> SPEVIGO	<input type="checkbox"/> 900 mg IV infusion over 90 minutes <input type="checkbox"/> Additional 900 mg IV infusion over 90 minutes one week after initial dose if flare symptoms persist	_____
<input type="checkbox"/> STELARA	Psoriasis Adult Subcutaneous <input type="checkbox"/> For patients <= 100 kg, 45 mg SC injection initially and 4 weeks later, followed by 45 mg every 12 weeks <input type="checkbox"/> For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks	_____
	Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose) <input type="checkbox"/> For patients <= 60 kg, 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks <input type="checkbox"/> For patients 60 kg – 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks <input type="checkbox"/> For patients >100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks	_____
	Psoriatic Arthritis Adult <input type="checkbox"/> 45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks <input type="checkbox"/> For patients with co-existent moderate-to-severe plaque psoriasis weighing >100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks	_____
<input type="checkbox"/> XOLAIR	<input type="checkbox"/> 150 or <input type="checkbox"/> 300 mg SC injection once every 4 weeks	_____
<input type="checkbox"/> IG	For Immunoglobulin therapy please refer to Immunoglobulin Form	_____
<input type="checkbox"/> OTHER		_____

By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature _____
 Dispense as Written _____
 Print Name _____
 Date _____

Prescriber's Signature _____
 Substitution Permitted _____
 Print Name _____
 Date _____

