

# Dermatology Referral Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION			
Patient Name:	Date of Birth:	Referral Date:	
Address:	City/State/Zip:		
Home Phone:	Cell Phone:	Work Phone:	
Secondary Contact:	Height:	Weight:	<input type="checkbox"/> Male <input type="checkbox"/> Female
Patient Diagnosis & ICD-10:			
Allergies:			

PROVIDER INFORMATION			
Physician Name:	Lic.#:	DEA #:	
Practice Name:	NPI#:		
Address:	City/State/Zip:		
Office Contact:	Phone:	Fax:	
Supervisory Physician (if applicable):			

PLEASE ATTACH	
<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical)	<input type="checkbox"/> TB lab results within last 12 months ( <i>Stelara, Simponi Aria, Ilumya &amp; Infliximabs only</i> )
<input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results	<input type="checkbox"/> HBV lab results within last 12 months ( <i>Infliximabs &amp; Simponi Aria only</i> )
<input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates)	<input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guidelines

NURSING & LAB ORDERS	
<b>Nurse Orders:</b> Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.	
<b>Flush Orders:</b> NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - <input type="checkbox"/> 10units/mL ---OR--- <input type="checkbox"/> 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line	
<b>Lab Orders:</b> <b>Lab Date &amp; Frequency:</b>	

PRESCRIPTION ORDERS			
<b>Anaphylaxis Kit:</b>	<input type="checkbox"/> Epinephrine 0.3mg IM as needed	<input type="checkbox"/> Solu-cortef 250mg-500mg IV as needed	<input type="checkbox"/> Solu-Medrol 60mg - 125mg IV as needed
(Check all that apply)	<input type="checkbox"/> Diphenhydramine _____ mg IV as needed	<input type="checkbox"/> NS Hydration 500 ml IV over 30 minutes as needed	<input type="checkbox"/> Other _____
<b>Pre-Medications:</b>	<input type="checkbox"/> Acetaminophen _____ mg PO _____ minutes prior to infusion	<input type="checkbox"/> Solu-Medrol _____ mg IV _____ minutes prior to infusion	
(Check all that apply)	<input type="checkbox"/> Diphenhydramine _____ mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV infusion _____ minutes prior to infusion	<input type="checkbox"/> Other _____	
<b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No If No, when was last dose given? _____ When is patient due for next dose? _____		
<input type="checkbox"/> ILUMYA	100mg SC injection at 0 and 4 weeks then every 12 weeks	_____
<input type="checkbox"/> INFlixIMAB <input type="checkbox"/> Avsola <input type="checkbox"/> Inflectra <input type="checkbox"/> Remicade <input type="checkbox"/> Renflexis	<input type="checkbox"/> <b>Induction:</b> _____ mg/kg or _____ mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours at weeks 0, 2, and 6	NONE
	<input type="checkbox"/> <b>Maintenance:</b> _____ mg/kg _____ mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours every _____ weeks <i>(Note: Round to nearest 100mg for Medicaid patients)</i> If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.	_____
<input type="checkbox"/> SIMPONI ARIA	2 mg/kg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter	_____
<input type="checkbox"/> SPEVIGO	<input type="checkbox"/> 900 mg IV infusion over 90 minutes <input type="checkbox"/> Additional 900 mg IV infusion over 90 minutes one week after initial dose if flare symptoms persist	_____
<input type="checkbox"/> STELARA	<b>Psoriasis Adult Subcutaneous</b> <input type="checkbox"/> For patients <= 100 kg, 45 mg SC injection initially and 4 weeks later, followed by 45 mg every 12 weeks <input type="checkbox"/> For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks	_____
	<b>Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose)</b> <input type="checkbox"/> For patients <= 60 kg, 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks <input type="checkbox"/> For patients 60 kg – 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks <input type="checkbox"/> For patients >100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks	_____
	<b>Psoriatic Arthritis Adult</b> <input type="checkbox"/> 45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks <input type="checkbox"/> For patients with co-existent moderate-to-severe plaque psoriasis weighing >100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks	_____
<input type="checkbox"/> XOLAIR	<input type="checkbox"/> 150 or <input type="checkbox"/> 300 mg SC injection once every 4 weeks	_____
<input type="checkbox"/> IG	<b>For Immunoglobulin therapy please refer to Immunoglobulin Form</b>	_____
<input type="checkbox"/> OTHER		_____

*By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.*

Prescriber's Signature \_\_\_\_\_  
Dispense as Written

Print Name \_\_\_\_\_ Date \_\_\_\_\_

Prescriber's Signature \_\_\_\_\_  
Substitution Permitted

Print Name \_\_\_\_\_ Date \_\_\_\_\_

