Dermatology Referral Form

Fax Completed Form To: 877-418-4495 Phone: 877-418-4114



PATIENT INFORMATION							
Patient Name:		Date of Birth:	TINIONMATION		Referral Date:		
Address:		Date of Dirtif.		City/State/Zip			
Home Phone:		Cell Phone:		City/State/Zip	Work Phone:		
Secondary Contact:		Height:	Weight:		Male	Female	
Patient Diagnosis & ICD	10:						
Allergies:							
PROVIDER INFORMATION							
Physician Name:		Lic.#:		DEA #:			
Practice Name:				NPI#:			
Address:				City/State/Zip	:		
Office Contact:			Phone:		Fax:		
Supervisory Physician (if applicable):							
PLEASE ATTACH							
Patient demographics & front/back copy of all insurance cards (prescription & medical) TB lab results within last 12 months (Stelara, Simponi Aria, Ilumya & Infliximabs only)							
Recent office visit notes, history & physical, lab & pertinent procedure results HBV lab results within last 12 months (Infliximabs & Simponi Aria only)							
Current medication list & list of prior medications tried and failed (with dates) Letter of medical necessity if drug dosing or indication is outside of FDA guidelines							
NURSING & LAB ORDERS							
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.							
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL -3-5mL flush after post-infusion NS flush if indicated to maintain line							
Lab Orders: Lab Date & Frequency:							
PRESCRIPTION ORDERS							
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed							
(Check all that apply) Diphenhydramine mg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other							
Pre-Medications: Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IVminutes prior to infusion							
(Check all that apply) Diphenhydraminemg POOR IV infusionminutes prior to infusion Other							
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary							
PRODUCT PRESCRIPTION INFORMATION REFILLS							
Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose?							
ILUMYA	100mg SC injection at 0 and 4 weeks then every 12 weeks						
INFLIXIMAB	Induction:mg/kg ormg IV infusion via gravityOR pump over at least 2 hours at weeks 0, 2, and 6						NONE
Avsola	Maintenance:mg/kgmg IV infusion via gravityOR pump over at least 2 hours everyweeks						
Inflectra	(Note: Round to nearest 100mg for Medicaid patients)						
Remicade	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.						
Renflexis SIMPONI ARIA	2 mg/kg IV infusion via gravityOR pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter						
SPEVIGO	900 mg IV infusion over 90 minutes Additional 900 mg IV infusion over 90 minutes one week after initial dose if flare symptoms persist						
JI LVIGO	Psoriasis Adult Subcutaneous	Additional 200 mg W imas	Sion over 70 minutes one week	tarter irritiar do	se ii iiaie syirip	torns persist	
STELARA	For patients <= 100 kg, 45 mg SC injection	on initially and 4 weeks la	ter, followed by 45 mg every 1	2 weeks			
	For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks						
	Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose)						
	For patients <= 60 kg, 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks						
	For patients 60 kg — 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks For patients > 100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks						
	Psoriatic Arthritis Adult						
	45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks						
	For patients with co-existent moderate-to-severe plaque psoriasis weighing > 100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks						
XOLAIR	150 or 300 mg SC injection once every 4 weeks						
IG	For Immunoglobulin therapy please refer to Immunoglobulin Form						
OTHER							
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.							

Prescriber's Signature

<u>Dispense as Written</u>

Print Name

Date

Prescriber's Signature Substitution Permitted **Print Name**

Date





