Gastroenterology Referral Form





Fax Completed Form To:	Phone

PATIENT INFORMATION							
Patient Name:		Date of Birth: Referral Date:					
Address:				City/State/Zip	re/Zip:		
Home Phone:		Cell Phone:		Work Phone:			
Secondary Contact:		Height:	Weight:		☐ Male ☐ Female		
Patient Diagnosis & ICD-10:							
Allergies:							
PROVIDER INFORMATION PROVIDER INFORMATION							
Physician Name:		Lic.#:		DEA #:			
Practice Name:				NPI#:			
Address:				City/State/Zip:			
Office Contact:		Phone:		Fax:			
Supervisory Physician (i	fapplicable):						
PLEASE ATTACH							
☐ Patient demographics & front/back copy of all insurance cards (prescription & medical) ☐ Recent office visit notes, history & physical, lab & pertinent procedure results ☐ Current medication list & list of prior medications tried and failed (with dates) ☐ Line access documentation/verification if applicable ☐ Vaccine status (any vaccination) and documentation of any recent vaccinations ☐ IB lab results within last 12 months ☐ HBV lab results within last 12 months ☐ Liver enzymes lab results (Skyrizi only) ☐ Bilirubin levels (Skyrizi only) ☐ Letter of medical necessity if drug dosing on the procedure results ☐ Letter of medical necessity if drug dosing on the procedure results		·	25				
		NURSIN	NG & LAB ORDERS				
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - Ununits/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency:							
		PRESC	RIPTION ORDERS				
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed							
(Check all that apply) Diphenhydraminemg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other Pre-Medications: Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IVminutes prior to infusion (Check all that apply) Diphenhydraminemg POOR IVminutes prior to infusion Other							
` ''	plies for vascular access line care, drug adminis						
PRODUCT			ION INFORMATION			REFILLS	
Is this a first dose?	Yes 🔲 No If No, when was last dose giver		_ When is patient due for next				
□ ENTYVIO -	☐ Induction: 300mg IV infusion over 30 minutes at week 0, 2, and 6			NONE			
	☐ Maintenance: 300mg IV infusion over 30 minutes every weeksOR Prefilled Pen 108mg SC every 2 weeks starting at week 6					2 pens, 13 refills	
☐ INFLIXIMAB ☐ Avsola ☐ Inflectra	□ Induction: mg/kg ormg IV infusion via □ gravity OR □ pump over at least 2 hours at weeks 0, 2, and 6				hours at weeks 0, 2, and 6	NONE	
	☐ Maintenance :mg/kgmg/V infusion via ☐ gravity OR ☐ pump over at least 2 hours everyweeks (Note: Round to nearest 100mg for Medicaid patients)						
☐ Remicade ☐ Renflexis	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.						
☐ Induction: 300mg IV infusion via ☐ gravityOR ☐ pump over 30 minutes at week 0, 4, and 8					NONE		
	☐ Maintenance: 200mg SC injection (given as two consecutive injections of 100 mg each) at Week 12, and every 4 weeks thereafter						
□ SKYRIZI -	☐ Induction (Crohn's): 600mg IV infusion via ☐ gravityOR ☐ pump over one hour at week 0, 4, and 8 ☐ Maintenance: ☐ 180mg or ☐ 360mg SC injection at Week 12, and every 8 weeks thereafter			NONE			
	□ Induction (UC): 1200mg IV infusion via □ gravityOR □ pump over two hours at week 0, 4, and 8 □ Maintenance: □ 180mg or □ 360mg SC injection at Week 12, and every 8 weeks thereafter				NONE 		
□ STELARA	Induction (Adult Dosing -Based on body weight of patient at time of dosing): ☐ For patients 55kg or less administer 260mg IV infusion via ☐ gravityOR ☐ pump over at least 1 hour x 1 dose ☐ For patients more than 55kg to 85kg administer 390mg IV infusion via ☐ gravityOR ☐ pump over at least 1 hour x 1 dose ☐ For patients more than 85kg administer 520mg IV infusion via ☐ gravityOR ☐ pump over at least 1 hour x 1 dose ☐ How the second of the light of the second of the				NONE		
☐ TREMFYA	□ Maintenance: 90mg SubQ injectionweeks after induction and everyweeks thereafter			NONE			
	☐ Induction: 200mg IV infusion on weeks 0, 4, and 8 ☐ Maintenance: 100mg SubQ injection every 8 weeks beginning at week 16				NONE		
	Maintenance: 100mg SubQ injection every 8 weeks beginning at week 16 Maintenance: 200mg SubQ injection every 4 weeks beginning at week 12						
□ OTHER						NONE 	
By signing this fo	orm and utilizing our services, you are authoriz	ring Amerita to serve as yo	our prior authorization design	ated agent in a	lealing with medical and prescription ins	urance companies.	

Prescriber's Signature <u>Dispense</u> as Written

Print Name

Date

Prescriber's Signature Substitution Permitted

Print Name

Date





