Gastroenterology Referral Form

Fax Completed Form To:

Phone:



PATIENT INFORMATION						
Patient Name:		Date of Birth:			Referral Date:	
Address:			City/State/Zip:			
Home Phone: Cell Phone:		Cell Phone:		1	Work Phone:	
Secondary Contact: Height:		Weight:		☐ Male ☐ Female		
Patient Diagnosis & ICD-10:						
Allergies:						
PROVIDER INFORMATION PROVIDED IN THE PROVIDED						
Physician Name: Lic.#:				DEA#:		
Practice Name: NPI#:						
Address:			City/State/Zip:			
Office Contact: Phone:			Fax:			
Supervisory Physician (if applicable):						
PLEASE ATTACH						
 □ Patient demographics & front/back copy of all insurance cards (prescription & medical) □ Recent office visit notes, history & physical, lab & pertinent procedure results □ Current medication list & list of prior medications tried and failed (with dates) □ Line access documentation/verification if applicable □ Vaccine status (any vaccination) and documentation of any recent vaccinations □ B lab results within last 12 months □ Liver enzymes lab results (Skyrizi only) □ Bilirubin levels (Skyrizi only) □ Letter of medical necessity if drug dosing or indication is outside of FDA guideline 						S
NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency:						
PRESCRIPTION ORDERS						
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed (Check all that apply) Diphenhydramine mg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other						
Pre-Medications: ☐ Acetaminophen mg PO minutes prior to infusion ☐ Solu-Medrol mg IV minutes prior to infusion (Check all that apply) ☐ Diphenhydramine mg ☐ PO IV minutes prior to infusion ☐ Other						
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary						
PRODUCT PRESCRIPTION INFORMATION REFILLS						
Is this a first dose?						
□ ENTYVIO -	☐ Induction: 300mg IV infusion over 30 minutes at week 0, 2, and 6					NONE
	☐ Maintenance: 300mg IV infusion over 30 minutes every weeksOR Prefilled Pen 108mg SC every 2 weeks starting at week 6					2 pens, 13 refills
☐ INFLIXIMAB ☐ Avsola ☐ Inflectra ☐ Remicade	□ Induction:mg/kg ormg IV infusion via □ gravityOR□ pump over at least 2 hours at weeks 0, 2, and 6					NONE
	☐ Maintenance :mg/kgmg/V infusion via ☐ gravity OR ☐ pump over at least 2 hours everyweeks (Note: Round to nearest 100mg for Medicaid patients)					
□Renflexis	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.					
□ омуон	☐ Induction: 300mg IV infusion via ☐ gravityOR ☐ pump over 30 minutes at week 0, 4, and 8					NONE
	☐ Maintenance: 200 mg SC injection (given as two consecutive injections of 100 mg each) at Week 12, and every 4 weeks thereafter					
□ SKYRIZI	☐ Induction (Crohn's): 600mg IV infusion via ☐ gravity0R ☐ pump over one hour at week 0, 4, and 8 ☐ Maintenance: ☐ 180mg or ☐ 360mg SC injection at Week 12, and every 8 weeks thereafter					NONE
	☐ Induction (UC): 1200mg IV infusion via ☐ gravityOR ☐ pump over two hours at week 0, 4, and 8 ☐ Maintenance: ☐ 180mg or ☐ 360mg SC injection at Week 12, and every 8 weeks thereafter					NONE
☐ STELARA	Induction (Adult Dosing -Based on body weight of patient at time of dosing): ☐ For patients 55kg or less administer 260mg IV infusion via ☐ gravity OR ☐ pump over at least 1 hour x 1 dose ☐ For patients more than 55kg to 85kg administer 390mg IV infusion via ☐ gravity OR ☐ pump over at least 1 hour x 1 dose ☐ For patients more than 85kg administer 520mg IV infusion via ☐ gravity OR ☐ pump over at least 1 hour x 1 dose ☐ Maintenance: 90mg SubQ injection weeks after induction and every weeks thereafter					NONE
☐ TREMFYA			uucuon anu everyV	אירבעי ווובובמונה		NONE
	☐ Induction: 200mg IV infusion on weeks 0, 4, and 8 ☐ Maintenance: 100mg SubQ injection every 8 weeks beginning at week 16					NONE
	☐ Maintenance: 200mg SubQ injection every 4 weeks beginning at week 12					NONE
□ OTHER	and addition and additional and additional a		and the state of t	-4-4 · · ·	allow and the second se	
By signing this fo	rm and utilizing our services, you are authorizi	ng Amerita to serve as yo	ur prior autnorization desiand	atea agent in de	aung with meaical and prescription ins	urance companies.

Prescriber's Signature <u>Dispense as Written</u>

Print Name

Date

Prescriber's Signature Substitution Permitted

Print Name

Date



