## Krystexxa® Order Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION						
Patient Name:	Date of Birth:	Referral Date:				
Address:		City/State/Zip:				
Home Phone:	Cell Phone:				k Phone:	
Secondary Contact:	Height:	Weig	ght:		Male 🛛 Female	
Patient Diagnosis & ICD-10:						
PROVIDER INFORMATION						
Physician Name:	Lic.#:			DEA #:		
Practice Name:	NPI#:					
Address: Office Contact:	City/State/Zip:					
	familicable):			Fax:		
Supervisory Physician (if applicable): PLEASE ATTACH						
				Evidence of patient on concurrent immunomodulation therapy such as: methotrexate, mycophenolate, leflunomide, azathioprine, or cyclosporine (Evidence supports the		
					imunomodulator in improving th	
Current medication	therapy; consider adding an immunomodulator if clinically appropriate.)					
G6PD deficiency results			□ Baseline serum Uric Acid lab results			
□ Verification that patient has discontinued or plans to discontinue oral urate lowering medications			□ Letter of medical necessity if drug dosing or indication is outside of FDA guidelines			
NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.						
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 🗆 10units/mLOR 🗆 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line						
Lab Orders:						
Lab Date & Frequency:						
PRESCRIPTION ORDERS						
Anaphylaxis Kit: 🛛 Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV infusion as needed Solu-Medrol 60mg - 125mg IV infusion as needed						
(Check all that apply) 🗆 Diphenhydramine mg IV infusion as needed 📄 NS Hydration 500 ml IV infusion over 30 minutes as needed 📄 Other						
Pre-Medications: Per prescribing information: Pre-medications of antihistamine, corticosteroid and analgesic is recommended						
(Check all that apply) 🗆 Acetaminophenmg POminutes prior to infusion 🖾 Solu-Medrolmg IV infusionminutes prior to infusion						
Diphenhydramine mg D - D OR D IV infusion minutes prior to infusion mg to infusion more spinor to infusion d ther						
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary						
PRODUCT			NFORMATIO			REFILLS
Is this a first dose? Yes No If No, when was last dose given?When is patient due for next dose?						
	□ 8mg IV infusion via □ gravity <b>OR</b> □ pump over at	least 2 hours eve	ery 2 weeks			
	□ After first infusion, patient to have sUA level performed within 48 hours prior to each infusion.					
□ Krystexxa	For KVO: NS 100mL via IV infusion over 1 hour.					
	If sUA is $\leq$ 6mg/dL, <b>proceed</b> .					
	If sUA is > 6mg/dL, <b>hold &amp; contact provider.</b>					
□ OTHER						
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.						

Prescriber's Signature <u>Dispense as Written</u> Date

Prescriber's Signature Substitution Permitted Print Name

Date

ACCREDITED



