

LEMTRADA® Referral Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION

Patient Name:	Date of Birth:	Referral Date:
Address:		City/State/Zip:
Home Phone:	Cell Phone:	Work Phone:
Secondary Contact:	Height: Weight:	<input type="checkbox"/> Male <input type="checkbox"/> Female
Patient Diagnosis & ICD-10:		
Allergies:		

PROVIDER INFORMATION

Physician Name:	Lic.#:	DEA #:
Practice Name:		NPI#:
Address:		City/State/Zip:
Office Contact:	Phone:	Fax:
Supervisory Physician (if applicable):		

MS CLINICAL DETAILS

Type of MS: Primary progressive multiple sclerosis (PPMS) ---OR--- Relapsing multiple sclerosis (RMS)
Ambulation status: Able to ambulate more than 5 meters Able to ambulate without aid or rest for at least 100 meters
Relapse details: Two or more relapses within the previous two years One relapse within the previous year

PLEASE ATTACH

<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical)	<input type="checkbox"/> CBC with differential, Serum creatinine levels, urinalysis with cell counts, urine protein to creatinine ratio thyroid function tests
<input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results	<input type="checkbox"/> Pregnancy test results (if applicable)
<input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates)	<input type="checkbox"/> Vaccine status (any vaccination) and documentation of any recent vaccinations
<input type="checkbox"/> Line access documentation/verification if applicable	<input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guidelines

NURSING & LAB ORDERS

Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line
Oxygen: Give O₂ at 2L/M per nasal cannula as needed
Lab Orders: **Lab Date & Frequency:**

SUPPLY ORDERS

Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
---------	--------------------------	---------

<p>Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No If No, when was last dose given? _____ When is patient due for next dose? _____</p>	<p><input type="checkbox"/> LEMTRADA</p> <p><input type="checkbox"/> Pre Meds: Hydroxyzine HCl 50mg po prior to start of infusion and every 6 hours prn #25 Acyclovir 200mg po BID for a minimum of 2 months or until CD4+ count is > or = to 200 cells per microliter, whichever occurs later #60 Refill: #1 Cetirizine 10mg po prior to Lemtrada infusion Ondansetron 4mg po prn #25 Promethazine 25mg po prn #25 Famotidine 20mg prior to start of alemtuzumab infusion Acetaminophen 1000mg po prior to start of Lemtrada infusion and q6h prn Other: _____ Note – If needed, please send pain prescription to retail pharmacy</p> <p><input type="checkbox"/> Pre Infusion: Solu-Medrol 1000mg IV infusion in 500mL of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only Normal saline 0.9% 500ml IV prior to Lemtrada infusion on days 4 and 5</p> <p><input type="checkbox"/> Initial Course: 12mg/day IV infusion via <input type="checkbox"/> pump ---OR--- <input type="checkbox"/> gravity over 4 hours for 5 consecutive days</p> <p><input type="checkbox"/> Subsequent Course: 12mg/day IV infusion via <input type="checkbox"/> pump ---OR--- <input type="checkbox"/> gravity over 4 hours for 3 consecutive days *To start at least 12 months after previous dose*</p> <p><input type="checkbox"/> Post Meds: Normal saline 0.9% 500mL IV infusion over 1 hour post Lemtrada infusion</p>	_____
<p><input type="checkbox"/> ANAPHYLAXIS / SIDE EFFECT ORDERS</p>	<p><input type="checkbox"/> Ondansetron 4-8mg in 50-100mL 0.9% NaCl IV infusion over 15 minutes prn nausea</p> <p><input type="checkbox"/> Epinephrine 1:1000 0.5-1mL IVP prn angioedema/hypotension/bronchospasm/generalized urticaria</p> <p><input type="checkbox"/> Ketorolac: 30mg IVP over 3-5 minute</p> <p><input type="checkbox"/> Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx 15 mins prn pruritis/rash</p>	_____
<p><input type="checkbox"/> OTHER</p>		_____

By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature	Print Name	Date	Prescriber's Signature	Print Name	Date
Dispense as Written			Substitution Permitted		