

Multiple Sclerosis Referral Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION		
Patient Name:	Date of Birth:	Referral Date:
Address:	City/State/Zip:	
Home Phone:	Cell Phone:	Work Phone:
Secondary Contact:	Height:	Weight: <input type="checkbox"/> Male <input type="checkbox"/> Female
Patient Diagnosis & ICD-10:		
Allergies:		

PROVIDER INFORMATION		
Physician Name:	Lic.#:	DEA #:
Practice Name:	NPI#:	
Address:	City/State/Zip:	
Office Contact:	Phone:	Fax:
Supervisory Physician (if applicable):		

MS CLINICAL DETAILS
Type of MS: <input type="checkbox"/> Primary progressive multiple sclerosis (PPMS) ---OR--- <input type="checkbox"/> Relapsing multiple sclerosis (RMS) Ambulation status: <input type="checkbox"/> Able to ambulate more than 5 meters <input type="checkbox"/> Able to ambulate without aid or rest for at least 100 meters Relapse details: <input type="checkbox"/> Two or more relapses within the previous two years <input type="checkbox"/> One relapse within the previous year

PLEASE ATTACH	
<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical) <input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results <input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates) <input type="checkbox"/> Line access documentation/verification if applicable	<input type="checkbox"/> Quantitative serum Immunoglobulin lab results (<i>Ocrevus only</i>) <input type="checkbox"/> Vaccine status (any vaccination) and documentation of any recent vaccinations <input type="checkbox"/> HBV lab results within last 12 months (<i>Ocrevus only</i>) <input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guideline

NURSING & LAB ORDERS
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - <input type="checkbox"/> 10units/mL ---OR--- <input type="checkbox"/> 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency:

PRESCRIPTION ORDERS
Anaphylaxis Kit: <input type="checkbox"/> Epinephrine 0.3mg IM as needed <input type="checkbox"/> Solu-cortef 250mg-500mg IV infusion as needed <input type="checkbox"/> Solu-Medrol 60mg - 125mg IV infusion as needed (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg IV infusion as needed <input type="checkbox"/> NS Hydration 500 ml IV infusion over 30 minutes as needed <input type="checkbox"/> Other
Pre-Medications: <input type="checkbox"/> Acetaminophen _____ mg PO _____ minutes prior to infusion <input type="checkbox"/> Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV infusion _____ minutes prior to infusion <input type="checkbox"/> Other
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No If No, when was last dose given? _____ When is patient due for next dose? _____		
<input type="checkbox"/> OCREVUS	<input type="checkbox"/> Induction: 300mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2.5 hours followed 2 weeks later by 300mg IV infusion over at least 2.5 hours <input type="checkbox"/> Maintenance: 600mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 3.5 hours every 6 months (if no prior serious infusion reactions, may administer over at least 2 hours) Post Infusion: Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion (Per PI, Corticosteroid and antihistamine required for pre-medication, refer to section above)	NONE
<input type="checkbox"/> TYSABRI	300mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over one hour every 4 weeks Post Infusion: Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion	NONE
<input type="checkbox"/> IG	For Immunoglobulin therapy please refer to Immunoglobulin Form	
<input type="checkbox"/> LEMTRADA	For Lemtrada therapy please refer to Lemtrada Form	
<input type="checkbox"/> OTHER		

By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature	Print Name	Date	Prescriber's Signature	Print Name	Date
Dispense as Written			Substitution Permitted		