Neurology Order Form Fax Completed Form To:



Phone:

rax completed i				
	PATIENT IN	NFORMATION		
Patient Name: Date of Birth:			Referral Date:	
Address: City/State/Zip:				
Home Phone:	Cell Phone:		Work Phone:	
Secondary Contact:		Weight:	☐ Male ☐ Female	
Patient Diagnosis & ICD-10:				
Allergies:				
PROVIDER INFORMATION				
Physician Name:	Lic.#:	DEA #:	<u>: </u>	
Practice Name:		NPI#:		
Address:	·	City/St	tate/Zip:	
Office Contact:	Phone:		Fax:	
Supervisory Physician (if applicable):				
PLEASE ATTACH				
☐ Patient demographics & front/back copy of all insurance cards (prescription & medical) ☐ Vaccine status (any vaccination) and documentation of any recent vaccinations				
☐ Recent office visit notes, history & physical, lab & pertinent procedure results ☐ HBV lab results within last 12 months (<i>Uplizna only</i>)				
☐ Current medication list & list of prior medications tried and failed (with dates) ☐ Date of diagnosis, current FVC%, ALSFRS-R score, and JourneyMate form (Radicava only)				
☐ Line access documentation/verification if applicable ☐ Anti-acetylcholine receptor (AChR) antibody positive results (<i>Vyvgart</i>)				
☐ Quantitative serum Immunoglobulin lab results (<i>Uplizna only</i>) ☐ Letter of medical necessity if drug dosing or indication is outside of FDA guidelines				
☐ TB lab results within last 12 months (<i>Uplizna only</i>)				
NURSING & LAB ORDERS				
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.				
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL OR 110units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line				
Lab Orders: Lab Date & Frequency:				
PRESCRIPTION ORDERS				
Anaphylaxis Kit: ☐ Epinephrine 0.3mg IM as needed ☐ Solu-cortef 250mg-500mg IV as needed ☐ Solu-Medrol 60mg - 125mg IV as needed				
(Check all that apply)				
Pre-Medications: ☐ Acetaminophenmg PO minutes prior to infusion ☐ Solu-Medrolmg IVminutes prior to infusion				
(Check all that apply)				
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary				
PRODUCT PRESCRIPTION INFORMATION RE				REFILLS
Is this a first dose?				
☐ Induction: 700mg IV infusion via ☐ gravityOR ☐ pump over 30 minutes every 4 weeks x 3 doses				NONE
□ KINSULA				NONE
	□ Maintenance: 1400mg IV infusion via □ gravityOR □ pump over 30 minutes every 4 weeks			
	☐ If missed dose, administer the same dose as soon as possible and continue every 4 weeks			
☐ RADICAVA	□ Induction: 60mg IV infusion via □ gravityOR □ pump over 1 hour daily for 14 days followed by 14 day drug-free period			NONE
	☐ Maintenance : 60mg IV infusion via ☐ gravityOR ☐ pump over 1 hour daily for 10 days out of 14 day period followed by 14 day drug-free periods			
	□ Induction : 300mg IV infusion via □ gravityOR □ pump over approximately 90 minutes at 0 and 2 weeks and CBC lab testing everymonths			NONE
LJ UPLIZNA -	□ Maintenance : (starting 6 months from first infusion) 300mg IV infusion via □ gravity 0R □ pump over approximately 90 minutes every 6 months			HOHE
□ VYEPTI	□ 100mg IV infusion via □ gravity OR □ pump over approximately 30 minutes every 12 weeks			
	□300mg IV infusion via □ gravityOR □ pump over approximately 30 minutes every 12 weeks □ □ □ □			
□ VYVGART	10mg/kg IV infusion via ☐ gravity OR pump over at least	1 hour once every week for 4 we	eks	
	*Up to max of 1200mg for patient weight of 120kg+ (Total volume is 125ml in NS solution)			
	Administer additional treatment cycles \square every 50 days OR \square Prescriber to evaluate treatment cycle frequency after completion of initial treatment cycle			
	According to the Package Insert: Administer subsequent treatment cycles based on clinical evaluation; the safety of initiating subsequent cycles sooner than 50 days			
	from the start of the previous treatment cycle has not been established.			
	1,008mg/11,200 units subcutaneous injection over approximately 30 to 90 seconds in cycles of once weekly injections for 4 weeks			
☐ VYVGART HYTRULO	Administer additional treatment cycles every 50 days OR Prescriber to evaluate treatment cycle frequency after completion of initial treatment cycle			
VIVOAMIIIIMOLO	According to the Package Insert: Administer subsequent treatment cycles based on clinical evaluation; the safety of initiating subsequent cycles sooner than 50 days			
	from the start of the previous treatment cycle has not been established.			
□ IG	Refer to Immunoglobulin Form			
☐ SOLIRIS/ULTOMIRIS	Refer to Soliris or Ultomiris Order Form			
☐ OTHER				NONE
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.				

Prescriber's Signature Dispense as Written

Print Name

Date

Prescriber's Signature **Substitution Permitted** Print Name

Date





