## **Pulmonary** Referral Form





Fax Completed Form To:

P	hc	e:

PATIENT INFORMATION								
Patient Name:	Date of Birth:		Referral Date:					
Address:	L			City/State/Zip:				
Home Phone:		Cell Phone:	Weisk		ork Phone:			
Secondary Contact: Patient Diagnosis & ICD	10.	Height:	Weight:	L	] Male ☐ Female			
Allergies:	-10.							
PROVIDER INFORMATION								
Physician Name:		Lic.#:		DEA #:				
Practice Name:		NPI#:						
Address:		City/State/Zip:						
Office Contact:		Fax:						
Supervisory Physician (i	f applicable):							
PLEASE ATTACH								
☐ Patient demographi	mographics & front/back copy of all insurance cards (prescription & medical)							
	otes, history & physical, lab & pertinent proc		☐ Alpha-1 antitrypsin leve		sia only)			
	list & list of prior medications tried and faile	d (with dates)	☐ FEV1 score (Aralast and					
	□ Documentation on phenotype (Aralast and Glassia only) □ Current Smoker? □ Yes □ No (Aralast and Glassia only)							
	Aralast and Glassia only)		☐ Line access documentat		•	i		
☐ CT scan results (Aralo	•		Letter of medical necess	ity it arug aosing or	indication is outside of FDA guidel	ines		
iga ievei (Araiast ar.	ia diassia only)							
NURSING & LAB ORDERS								
	provide assessment, teaching, lab draws, m			-				
	$\%$ - 5-10mL flush pre and post infusion and $\alpha$	as needed <i>Heparin</i> - ∟ 1		ınits/mL - 3-5mL flu	ish after post-infusion NS flush if ir	idicated to maintain line		
Lab Orders:			Lab Date & Frequency:					
A 1 1 ' 10'			RIPTION ORDERS		C   M	· 1.1		
Anaphylaxis Kit: (Check all that apply)	☐ Epinephrine 0.3mg IM as needed☐ Diphenhydraminemg I	□ Solu-cortef 250mg-5 V infusion as needed	Solomg IV Infusion as needed  NS Hydration 500 ml IV		Solu-Medrol 60mg - 125mg IV infi	usion as needed  Other		
Pre-Medications:	☐ Acetaminophenmg PO		· · · · · · · · · · · · · · · · · · ·			□ otner		
Pre-Medications:       □ Acetaminophenmg POminutes prior to infusion       □ Solu-Medrolmg IVminutes prior to infusion         (Check all that apply)       □ Diphenhydraminemg □ POOR □ IV infusionminutes prior to infusion □ Other								
11./:	plies for vascular access line care, drug admi							
PRODUCT	-		ION INFORMATIO			REFILLS		
Is this a first dose?  No If No, when was last dose given? When is patient due for next dose?								
☐ ARALAST	60mg/kg IV infusion via ☐ gravity <b>OR</b> ☐ pump weekly over approximately 15 minutes  *Administer at a rate not to exceed 0.2 mL/kg body weight per minute **Acceptable allotment +/- 10% based on vial lot/batch							
☐ CINQAIR	3mg/kg IV infusion via □gravity <b>OR</b> □ pump once every 4 weeks over 20-50 minutes							
☐ FASENRA	☐ Induction: 30mg SubQ injection every 4 weeks for the first 3 doses ☐ Maintenance: 30mg SubQ injection once every 8 weeks					NONE		
	60mg/kg IV infusion via gravity <b>OR</b> pump once weekly over approximately 15 minutes							
☐ GLASSIA	*Administer at a rate not to exceed 0.2 mL/kg bo							
□ NUCALA	□ 100mg SubQ injection every 4 weeks □ 300mg SubQ injection every 4 weeks □							
☐ TEZSPIRE	210mg SubQ injection once every 4 weeks							
☐ XOLAIR	mg SubQ injection every							
□ OTHER								
By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.								
Prescriber's Signature <u>Dispense as Written</u>	Print Name	Date	Prescriber's Signa Substitution Peri		Print Name	Date		







