TEPEZZA® Referral Form





Fax Completed Form To: Phone:

| PATIENT INFORMATION | | | | | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|----------------------------------------|-----------------|-------------|-------------------------------|
| Patient Name: | Date of Birth: | | Referral Date: | | | |
| Address: | | | | City/State/Zip: | | |
| Home Phone: | | Cell Phone: | | | Work Phone: | |
| Secondary Contact: | | | Weight: | ☐ Male ☐ Female | | |
| Patient Diagnosis & ICD-10: | | | | | | |
| Allergies: | | | | | | |
| PROVIDER INFORMATION | | | | | | |
| Physician Name: | Lic.#: | | | DEA#: | | |
| Practice Name: | | | | NPI#: | | |
| Address: | Diama | | | City/State/Zip: | | |
| Office Contact: | Phone: | | | Fax: | | |
| Supervisory Physician (if applicable): PLEASE ATTACH | | | | | | |
| □ Patient demographics & front/back copy of all insurance cards (prescription & medical) □ Recent office visit notes, history & physical, lab & pertinent procedure results □ Current medication list & list of prior medications tried and failed (with dates) □ History of IBD documentation □ Diabetic documentation □ Prior treatments for TED: steroids, surgeries, or other treatments □ CAS score □ Thyroid lab results □ Notes detailing if mild or moderate TED □ Documentation of lid retraction of 2 or more millimeters or □ Documentation of proptosis of 3 millimeters or more □ Letter of medical necessity if drug dosing or indication is outside of FDA guideling a second course | | | | | | es or if patient is receiving |
| NURSING & LAB ORDERS | | | | | | |
| Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Routine/Standing Lab Orders: (attach if needed) | | | | | | |
| PRESCRIPTION ORDERS | | | | | | |
| Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV infusion as needed Solu-Medrol 60mg - 125mg IV infusion as needed Solu-Medrol 60mg - 125mg IV infusion as needed Other | | | | | | |
| Pre-Medications: □ Acetaminophenmg POminutes prior to infusion □ Solu-Medrolmg IV infusionminutes prior to infusion [Check all that apply] □ Diphenhydraminemg POOR □ IV infusionminutes prior to infusion □ Other | | | | | | |
| Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary | | | | | | |
| PRODUCT | PRESCRIPTION INFORMATION | | | | | REFILLS |
| Is this a first dose? | | | | | | |
| | □ INDUCTION: 10mg/kg IV infusion via □ gravityOR □ pump over 90 minutes for one time dose | | | | | NONE |
| ☐ TEPEZZA | ☐ MAINTENANCE: Maintenance: 20mg/kg IV infusion via ☐ gravityOR ☐ pump over 60 to 90 minutes every 3 weeks for 7 additional infusions | | | | | |
| | Administer the diluted solution intravenously over 90 minutes for the first two infusions. If well tolerated, the minimum time for subsequent infusions can be reduced to 60 minutes. If not well tolerated, the minimum time for subsequent infusions should remain at 90 minutes. | | | | | NONE |
| □ OTHER | | | | | | |
| By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies. | | | | | | |
| | | | | | | |
| Prescriber's Signature Dispense as Written | Print Name | Date | Prescriber's Signa Substitution Per | | Print Name | Date |





