

Gastroenterology Referral Form

Fax Completed Form To:

Phone:



Ivy Specialty Infusion
an amerita company

PATIENT INFORMATION			
Patient Name:		Date of Birth:	Referral Date:
Address:		City/State/Zip:	
Home Phone:	Cell Phone:	Work Phone:	
Secondary Contact:	Height:	Weight:	<input type="checkbox"/> Male <input type="checkbox"/> Female
Patient Diagnosis & ICD-10:			
Allergies:			
PROVIDER INFORMATION			
Physician Name:		Lic.#:	DEA #:
Practice Name:		NPI#:	
Address:		City/State/Zip:	
Office Contact:	Phone:		Fax:
Supervisory Physician (if applicable):			
PLEASE ATTACH			
<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical)		<input type="checkbox"/> TB lab results within last 12 months	
<input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results		<input type="checkbox"/> HBV lab results within last 12 months (<i>Infliximabs only</i>)	
<input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates)		<input type="checkbox"/> Liver enzymes lab results (<i>Skyrizi only</i>)	
<input type="checkbox"/> Line access documentation/verification if applicable		<input type="checkbox"/> Bilirubin levels (<i>Skyrizi only</i>)	
<input type="checkbox"/> Vaccine status (any vaccination) and documentation of any recent vaccinations		<input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guidelines	
NURSING & LAB ORDERS			
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.			
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - <input type="checkbox"/> 10units/mL ---OR--- <input type="checkbox"/> 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line			
Lab Orders: Lab Date & Frequency:			
PRESCRIPTION ORDERS			
Anaphylaxis Kit: <input type="checkbox"/> Epinephrine 0.3mg IM as needed <input type="checkbox"/> Solu-cortef 250mg-500mg IV as needed <input type="checkbox"/> Solu-Medrol 60mg - 125mg IV as needed (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg IV as needed <input type="checkbox"/> NS Hydration 500 ml IV over 30 minutes as needed <input type="checkbox"/> Other			
Pre-Medications: <input type="checkbox"/> Acetaminophen _____ mg PO _____ minutes prior to infusion <input type="checkbox"/> Solu-Medrol _____ mg IV _____ minutes prior to infusion (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV _____ minutes prior to infusion <input type="checkbox"/> Other			
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			
PRODUCT	PRESCRIPTION INFORMATION		REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No If No, when was last dose given? _____ When is patient due for next dose? _____			
<input type="checkbox"/> ENTYVIO	<input type="checkbox"/> Induction: 300mg IV infusion over 30 minutes at week 0, 2, and 6		NONE
	<input type="checkbox"/> Maintenance: 300mg IV infusion over 30 minutes every _____ weeks ---OR--- Prefilled Pen 108mg SC every 2 weeks starting at week 6		2 pens, 13 refills
<input type="checkbox"/> INFlixIMAB <input type="checkbox"/> Avsola <input type="checkbox"/> Inflectra <input type="checkbox"/> Remicade <input type="checkbox"/> Renflexis	<input type="checkbox"/> Induction: _____ mg/kg or _____ mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours at weeks 0, 2, and 6		NONE
	<input type="checkbox"/> Maintenance: _____ mg/kg _____ mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours every _____ weeks (Note: Round to nearest 100mg for Medicaid patients) If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.		_____
<input type="checkbox"/> OMVOH	<input type="checkbox"/> Induction (UC): 300mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 30 minutes at week 0, 4, and 8		NONE
	<input type="checkbox"/> Maintenance: 200mg SC injection, given as two consecutive injections of 100mg each at Week 12, and every 4 weeks thereafter		_____
<input type="checkbox"/> SKYRIZI	<input type="checkbox"/> Induction (Crohn's): 900mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 30 minutes at week 0, 4, and 8		NONE
	<input type="checkbox"/> Maintenance: 300mg SC injection, given as two consecutive injections of 100mg + 200mg (or 100mg x 3) at Week 12, and every 4 weeks thereafter		_____
	<input type="checkbox"/> Induction (UC): 600mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over one hour at week 0, 4, and 8		NONE
<input type="checkbox"/> STELARA	<input type="checkbox"/> Maintenance: <input type="checkbox"/> 180mg or <input type="checkbox"/> 360mg SC injection at Week 12, and every 8 weeks thereafter		_____
	<input type="checkbox"/> Induction (UC): 1200mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over two hours at week 0, 4, and 8		NONE
	<input type="checkbox"/> Maintenance: <input type="checkbox"/> 180mg or <input type="checkbox"/> 360mg SC injection at Week 12, and every 8 weeks thereafter		_____
<input type="checkbox"/> TREMFYA	Induction (Adult Dosing - Based on body weight of patient at time of dosing): <input type="checkbox"/> For patients 55kg or less administer 260mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 1 hour x 1 dose <input type="checkbox"/> For patients more than 55kg to 85kg administer 390mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 1 hour x 1 dose <input type="checkbox"/> For patients more than 85kg administer 520mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 1 hour x 1 dose		NONE
	<input type="checkbox"/> Maintenance: 90mg SubQ injection _____ weeks after induction and every _____ weeks thereafter		_____
<input type="checkbox"/> OTHER	<input type="checkbox"/> Induction: 200mg IV infusion on weeks 0, 4, and 8		NONE
	<input type="checkbox"/> Maintenance: 100mg SubQ injection every 8 weeks beginning at week 16 <input type="checkbox"/> Maintenance: 200mg SubQ injection every 4 weeks beginning at week 12		_____

By signing this form and utilizing our services, you are authorizing Amerita to assist with prior authorization requests acting as your pharmacy provider in dealing with medical and prescription insurance companies.

Prescriber's Signature
Dispense as Written

Print Name

Date _____

Prescriber's Signature
Substitution Permitted

Print Name

Date _____

©2025 Amerita Inc. All rights reserved.
AME IVY Gastroenterology Refer 6.25

ameritaiv.com

THIS FORM IS NOT VALID FOR USE IN THE STATE OF ARIZONA.

ACHC
ACCREDITED

PCAB ACCREDITED
Compounding Pharmacists