

# Multiple Sclerosis Referral Form

Fax Completed Form To:

Phone:

PATIENT INFORMATION		
Patient Name:	Date of Birth:	Referral Date:
Address:	City/State/Zip:	
Home Phone:	Cell Phone:	Work Phone:
Secondary Contact:	Height:	Weight: <input type="checkbox"/> Male <input type="checkbox"/> Female
Patient Diagnosis & ICD-10:		
Allergies:		
PROVIDER INFORMATION		
Physician Name:	Lic.#:	DEA #:
Practice Name:	NPI#:	
Address:	City/State/Zip:	
Office Contact:	Phone:	Fax:
Supervisory Physician (if applicable):		
MS CLINICAL DETAILS		
<b>Type of MS:</b> <input type="checkbox"/> Primary progressive multiple sclerosis (PPMS) ---OR--- <input type="checkbox"/> Relapsing multiple sclerosis (RMS)		
<b>Ambulation status:</b> <input type="checkbox"/> Able to ambulate more than 5 meters <input type="checkbox"/> Able to ambulate without aid or rest for at least 100 meters		
<b>Relapse details:</b> <input type="checkbox"/> Two or more relapses within the previous two years <input type="checkbox"/> One relapse within the previous year		
PLEASE ATTACH		
<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical)	<input type="checkbox"/> Quantitative serum Immunoglobulin lab results ( <i>Ocrevus only</i> )	
<input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results	<input type="checkbox"/> Vaccine status (any vaccination) and documentation of any recent vaccinations	
<input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates)	<input type="checkbox"/> HBV lab results within last 12 months ( <i>Ocrevus only</i> )	
<input type="checkbox"/> Line access documentation/verification if applicable	<input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guideline	
NURSING & LAB ORDERS		
<b>Nurse Orders:</b> Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.		
<b>Flush Orders:</b> NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - <input type="checkbox"/> 10units/mL ---OR--- <input type="checkbox"/> 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line		
<b>Lab Orders:</b> <span style="float: right;"><b>Lab Date &amp; Frequency:</b></span>		
PRESCRIPTION ORDERS		
<b>Anaphylaxis Kit:</b> <input type="checkbox"/> Epinephrine 0.3mg IM as needed <input type="checkbox"/> Solu-cortef 250mg-500mg IV infusion as needed <input type="checkbox"/> Solu-Medrol 60mg - 125mg IV infusion as needed (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg IV infusion as needed <input type="checkbox"/> NS Hydration 500 ml IV infusion over 30 minutes as needed <input type="checkbox"/> Other		
<b>Pre-Medications:</b> <input type="checkbox"/> Acetaminophen _____ mg PO _____ minutes prior to infusion <input type="checkbox"/> Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV infusion _____ minutes prior to infusion <input type="checkbox"/> Other		
<b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary		
PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No If No, when was last dose given? _____ When is patient due for next dose? _____		
<input type="checkbox"/> OCREVUS	<input type="checkbox"/> <b>Induction:</b> 300mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2.5 hours followed 2 weeks later by 300mg IV infusion over at least 2.5 hours <input type="checkbox"/> <b>Maintenance:</b> 600mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 3.5 hours every 6 months (if no prior serious infusion reactions, may administer over at least 2 hours) <b>Post Infusion:</b> Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion (Per PI, Corticosteroid and antihistamine required for pre-medication, refer to section above)	NONE  _____
<input type="checkbox"/> TYSABRI	300mg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over one hour every 4 weeks <b>Post Infusion:</b> Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion	NONE  _____
<input type="checkbox"/> IG	<b>For Immunoglobulin therapy please refer to Immunoglobulin Form</b>	
<input type="checkbox"/> LEMTRADA	<b>For Lemtrada therapy please refer to Lemtrada Form</b>	
<input type="checkbox"/> OTHER	_____	
<b>By signing this form and utilizing our services, you are authorizing Amerita to assist with prior authorization requests acting as your pharmacy provider in dealing with medical and prescription insurance companies.</b>		

Prescriber's Signature

Print Name

Date

Prescriber's Signature

Print Name

Date