

# TEPEZZA® Referral Form

Fax Completed Form To:

Phone:

PATIENT INFORMATION			
Patient Name:		Date of Birth:	
Address:		Referral Date:	
City/State/Zip:		Work Phone:	
Home Phone:	Cell Phone:	Height: Weight:	
Secondary Contact:	<input type="checkbox"/> Male <input type="checkbox"/> Female		
Patient Diagnosis & ICD-10:			
Allergies:			
PROVIDER INFORMATION			
Physician Name:		Lic.#:	
Practice Name:		DEA #:	
Address:		NPI#:	
City/State/Zip:		Fax:	
Office Contact:		Phone:	
Supervisory Physician (if applicable):			
PLEASE ATTACH			
<input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical) <input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results <input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates) <input type="checkbox"/> History of IBD documentation <input type="checkbox"/> Diabetic documentation <input type="checkbox"/> Prior treatments for TED: steroids, surgeries, or other treatments		<input type="checkbox"/> CAS score <input type="checkbox"/> Thyroid lab results <input type="checkbox"/> Notes detailing if mild or moderate TED <input type="checkbox"/> Documentation of lid retraction of 2 or more millimeters or <input type="checkbox"/> Documentation of proptosis of 3 millimeters or more <input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guidelines or if patient is receiving a second course	
NURSING & LAB ORDERS			
<b>Nurse Orders:</b> Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. <b>Flush Orders:</b> NaCl 0.9% - 5-10mL flush pre and post infusion and as needed   Heparin - <input type="checkbox"/> 10units/mL ---OR--- <input type="checkbox"/> 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line <b>Routine/Standing Lab Orders:</b> (attach if needed) <input type="checkbox"/> Blood glucose test every _____ infusion(s). <input type="checkbox"/> Pregnancy test prior to each infusion if childbearing age. <b>Lab Orders:</b> <b>Lab Date &amp; Frequency:</b>			
PRESCRIPTION ORDERS			
<b>Anaphylaxis Kit:</b> <input type="checkbox"/> Epinephrine 0.3mg IM as needed <input type="checkbox"/> Solu-cortef 250mg-500mg IV infusion as needed <input type="checkbox"/> Solu-Medrol 60mg - 125mg IV infusion as needed (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg IV infusion as needed <input type="checkbox"/> NS Hydration 500 ml IV infusion over 30 minutes as needed <input type="checkbox"/> Other			
<b>Pre-Medications:</b> <input type="checkbox"/> Acetaminophen _____ mg PO _____ minutes prior to infusion <input type="checkbox"/> Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion (Check all that apply) <input type="checkbox"/> Diphenhydramine _____ mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV infusion _____ minutes prior to infusion <input type="checkbox"/> Other			
<b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			
PRODUCT	PRESCRIPTION INFORMATION		REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> No   If No, when was last dose given? _____ When is patient due for next dose? _____			
<input type="checkbox"/> TEPEZZA	<input type="checkbox"/> <b>INDUCTION:</b> 10mg/kg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 90 minutes for one time dose		NONE
	<input type="checkbox"/> <b>MAINTENANCE:</b> Maintenance: 20mg/kg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 60 to 90 minutes every 3 weeks for 7 additional infusions <input type="checkbox"/> Administer the diluted solution intravenously over 90 minutes for the first two infusions. If well tolerated, the minimum time for subsequent infusions can be reduced to 60 minutes. If not well tolerated, the minimum time for subsequent infusions should remain at 90 minutes.		NONE
<input type="checkbox"/> OTHER			
<b>By signing this form and utilizing our services, you are authorizing Amerita to assist with prior authorization requests acting as your pharmacy provider in dealing with medical and prescription insurance companies.</b>			

Prescriber's Signature  
Dispense as Written

Print Name

Date

Prescriber's Signature  
Substitution Permitted

Print Name

Date