

Krystexxa® Order Form

Fax Completed Form To:

Phone:

PATIENT INFORMATION		
Patient Name:	Date of Birth:	Referral Date:
Address:		City/State/Zip:
Home Phone:	Cell Phone:	Work Phone:
Secondary Contact:	Height:	Weight: ☐ Male ☐ Female
Patient Diagnosis & ICD-10:		
Allergies:		
PROVIDER INFORMATION		
Physician Name:	Lic.#:	DEA #:
Practice Name:		NPI#:
Address:		City/State/Zip:
Office Contact:	Phone:	Fax:
Supervisory Physician (if applicable):		
PLEASE ATTACH		
☐ Patient demographics & front/back copy of all insurance cards (prescription & medical) ☐ Recent office visit notes, history & physical, lab & pertinent procedure results ☐ Current medication list & list of prior medications tried and failed (with dates) ☐ G6PD deficiency results ☐ Verification that patient has discontinued or plans to discontinue oral urate lowering medications ☐ Evidence of patient on concurrent immunomodulation therapy such as: methotrexate, mycophenolate, leflunomide, azathioprine, or cyclosporine (Evidence supports the combination of Krystexxa and an immunomodulator in improving the patient's response to therapy; consider adding an immunomodulator if clinically appropriate.) ☐ Baseline serum Uric Acid lab results ☐ Letter of medical necessity if drug dosing or indication is outside of FDA guidelines		
NURSING & LAB ORDERS		
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - ☐ 10units/mL ---OR--- ☐ 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency:		
PRESCRIPTION ORDERS		
Anaphylaxis Kit: ☐ Epinephrine 0.3mg IM as needed ☐ Solu-cortef 250mg-500mg IV infusion as needed ☐ Solu-Medrol 60mg - 125mg IV infusion as needed (Check all that apply) ☐ Diphenhydramine _____mg IV infusion as needed ☐ NS Hydration 500 ml IV infusion over 30 minutes as needed ☐ Other		
Pre-Medications: Per prescribing information: Pre-medications of antihistamine, corticosteroid and analgesic is recommended (Check all that apply) ☐ Acetaminophen _____mg PO _____minutes prior to infusion ☐ Solu-Medrol _____mg IV infusion _____minutes prior to infusion ☐ Diphenhydramine _____mg ☐ PO ---OR--- ☐ IV infusion _____minutes prior to infusion ☐ Other		
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary		
PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Is this a first dose? ☐ Yes ☐ No If No, when was last dose given? _____ When is patient due for next dose? _____		
☐ Krystexxa	☐ 8mg IV infusion via ☐ gravity ---OR--- ☐ pump over at least 2 hours every 2 weeks ☐ After first infusion, patient to have sUA level performed within 48 hours prior to each infusion. For KVO: NS 100mL via IV infusion over 1 hour. If sUA is ≤ 6mg/dL, proceed. If sUA is > 6mg/dL, hold & contact provider. ☐ Has a ready-to-use vial: 8 mg/50 mL (50 mL)	
☐ OTHER		

Prescriber's Signature
Dispense as Written

Print Name

Date

Prescriber's Signature
Substitution Permitted

Print Name

Date